

FISHERY RESEARCH



FEDERAL AID IN FISH RESTORATION
Job Completion Report, Project F-73-R-6
Subproject II: River and Stream Investigations
Study VI: Middle Fork Salmon River
Fisheries Investigations



SH
351
S77
T4863
1985

Russ Thurow
Senior Fishery Research Biologist
January, 1985

SH
351
S77
T4863
1985

TABLE OF CONTENTS

	<u>Page</u>
ABSTRACT	1
INTRODUCTION	3
DESCRIPTION OF STUDY AREA.	4
OBJECTIVES	6
RECOMMENDATIONS.	7
METHODS.	8
Spawning Area Surveys.	8
Juvenile Distribution and Abundance.	9
Middle Fork Salmon River.	10
Tributaries	10
Origin of Middle Fork Steelhead.	10
Adult Steelhead Movements.	11
Main Salmon River Sport Fishery.	11
RESULTS.	12
Biological Characteristics	12
Spawner Characteristics and Densities.	12
Brush Creek	14
Marble Creek.	14
Monumental Creek.	14
Sheep Creek	14
Sulphur Creek	19
Wilson Creek.	19
Yellowjacket Creek.	19
Juvenile Distribution and Abundance.	19
Middle Fork	19
Tributaries	29
Bear Valley Creek	39
Big Creek Tributaries	39
Brush Creek	40
Camas Creek	40
Loon Creek.	44
Marble Creek.	44
Papoose Creek	46
Sheep Creek	46
Sulphur Creek	46
Wilson Creek.	46
Life History and Movements	48
Origin of Middle Fork Steelhead.	51
Main Salmon River Sport Fishery.	51

TABLE OF CONTENTS (Continued)

	<u>Page</u>
Estimated Exploitation of Middle Fork Stocks.	51
Fall 1983 Section 4 Fishery	52
DISCUSSION	53
Spawning Behavior.	53
Establishment of Escapement Goals.	56
Restoration of Steelhead Sport Fishing Opportunities	62
Consideration for Future Sport Fishing	66
Status of Indigenous Chinook Salmon.	67
Status of Westslope Cutthroat Trout.	68
The Role of Tributaries	70
ACKNOWLEDGEMENTS	73
LITERATURE CITED	74
APPENDICES	80

LIST OF TABLES

Table 1. Fishes present in the Middle Fork Salmon River drainage	13
Table 2. Numbers and densities of juvenile steelhead observed in Middle Fork Salmon River transects, 1983	20
Table 3. Number of fish sampled by hook-and-line and snorkeling in the Middle Fork Salmon River, July-August, 1980-1983	26
Table 4. Age frequency of juvenile steelhead observed in twenty Middle Fork Salmon River transects, 1980-1983.	27
Table 5. Numbers of fish (non-steelhead) observed in Middle Fork River transects, August, 1983.	28
Table 6. Numbers and sizes of fish observed by snorkeling in Middle Fork Salmon River tributaries, July-September, 1983	30
Table 7. Number of fish sampled by hook-and-line and snorkeling in Middle Fork Salmon River tributaries, July-August, 1981-1983.	33

LIST OF TABLES (Continued)

	<u>Page</u>
Table 8. Fish observed by snorkeling in Middle Fork Salmon River tributaries, July-August, 1983	34
Table 9. Numbers of fish sampled by hook-and-line in Middle Fork Salmon River tributaries, July-August, 1983.	41
Table 10. Results of spring and fall surveys for adult steelhead in the Middle Fork Salmon River drainage, 1983	49
Table 11. Summary of steelhead angler creel data for Salmon River Section 4, October-November, 1983.	54
Table 12. Steelhead angler effort and catch in Salmon River Section 4 during similar intervals, 1981-1983.	55
Table 13. Juvenile steelhead densities observed in underseeded and fully seeded Idaho streams	59
Table 14. Required escapements to produce 350,000 smolts in the Middle Fork Salmon River drainage with varying egg-to-smolt survival rates	61

LIST OF FIGURES

Figure 1. Middle Fork Salmon River drainage, Idaho.	5
Figure 2. Percent of steelhead spawners and redds (combined) observed in Big, Camas, and Loon creeks during specified periods, 1968-1983.	15
Figure 3. Spawning ground survey map of Brush and Sheep creeks, Middle Fork Salmon River, Idaho, 1983	16
Figure 4. Spawning ground survey map of Marble Creek, Middle Fork Salmon River, Idaho, 1983.	17
Figure 5. Spawning ground survey map of Monumental Creek, Middle Fork Salmon River, Idaho, 1983.	18
Figure 6. Spawning ground survey map of Sulphur Creek, Middle Fork Salmon River, Idaho, 1983.	21
Figure 7. Spawning ground survey map of Wilson Creek, Middle Fork Salmon River, Idaho, 1983.	22

LIST OF FIGURES (Continued)

	<u>Page</u>
Figure 8. Spawning ground survey map of Yellowjacket Creek, Middle Fork Salmon River, Idaho, 1983	23
Figure 9. Number and density of juvenile steelhead observed in Middle Fork Salmon River transects, 1980-1983	24
Figure 10. Length frequency of juvenile steelhead and cutthroat trout caught by angling in the Middle Fork Salmon River, 1980-1983.	31
Figure 11. Length frequency of juvenile steelhead and cutthroat trout caught by angling in Middle Fork Salmon River tributaries, 1981-1983.	36
Figure 12. Length frequency of bull trout caught by angling in Middle Fork Salmon River tributaries, 1981-1983	38
Figure 13. Length frequency of juvenile steelhead caught in Yellowjacket Creek, July, 1983.	43
Figure 14. Length frequency of juvenile steelhead and cutthroat trout caught in Marble Creek, 1982-1983	45
Figure 15. Length frequency of juvenile steelhead-rainbow caught in Wilson Creek, above and below a barrier, July, 1983	47
Figure 16. Numbers of adult steelhead observed in the lower 800 m of the Middle Fork Salmon River, 1981-1983.	50
Figure 17. Estimated annual escapements of wild steelhead trout to the Middle Fork Salmon River, 1970-1983.	65
Figure 18. The percentage of cutthroat trout exceeding 300 mm from hook-and-line samples from the Middle Fork Salmon River, 1959-1983.	69
Figure 19. Cutthroat trout densities at various elevations in Middle Fork Salmon River tributaries, 1982-1983	71

LIST OF APPENDICES

Appendix A. Manual for steelhead spawning ground surveys.	81
Appendix B. Data collected during steelhead spawner surveys (ground and aerial) on Middle Fork Salmon River tributaries, April-May, 1983.	95

LIST OF APPENDICES (Continued)

	<u>Page</u>
Appendix C. Physical dimensions and characteristics of snorkeling transects in the Middle Fork Salmon River, 2-9 August 1983.	97
Appendix D. Length frequency of juvenile steelhead and cutthroat trout caught by angling in the Middle Fork Salmon River and tributaries, 1983	98
Appendix E. Mean dimensions of snorkeling transects (mean of 5) in the Middle Fork Salmon River tributary sections, 1983.	99

JOB COMPLETION REPORT

State of: Idaho Name: RIVER AND STREAM INVESTIGATIONS
Project: F-73-R-6 Title: Middle Fork Salmon River
Fisheries Investigations
Subproject: II
Study: VI
Period Covered: 1 March 1983 - 29 February 1984

ABSTRACT

Status of wild steelhead trout (Salmo gairdneri) was evaluated in 14 tributaries to the Middle Fork Salmon River from 1981 to 1983. Emphasis was directed at collecting biological data to assist future management of the drainage.

Most of the drainage is in a pristine wilderness. Exceptions occur in several tributaries where man-caused activities (livestock grazing, placer gold and other precious metal mining) have degraded habitat. The proposed acceleration of mining within several drainages has the potential to degrade additional habitat.

Electrophoretic analysis suggests that Middle Fork Salmon River steelhead trout are similar to other "B" stock summer steelhead populations sampled in the Snake River basin. The data also illustrate that separate sub-populations exist within the Middle Fork Salmon River drainage.

Adult escapements in the study period were not sufficient to seed the available spawning habitat. Spawning commenced in early April and was completed prior to peak runoff in early June. We estimated that 640 km of tributaries were accessible to steelhead trout.

Tributaries provide the principal rearing habitats for steelhead trout in the Middle Fork drainage. Densities of steelhead trout ranged from 0.2 to 10.0 fish per 100 m² and averaged 4.0 fish per 100 m². Tributary rearing areas were underseeded from 1981 to 1983.

A small number of tributaries produce a majority of the westslope cutthroat trout (Salmo clarki) in the Middle Fork Salmon River. Bull trout (Salvelinus confluentus) were usually sympatric with cutthroat trout. Cutthroat trout populations were observed above migration barriers in three streams.

Few adult steelhead trout ascended the Middle Fork in the fall. Middle Fork Salmon River steelhead trout which stage in the mainstem Salmon River were formerly susceptible to an intensive sport fishery. A differential harvest regulation based on a 57 mm dorsal fin height measurement was instituted in the fall 1982. This regulation has allowed anglers to harvest a maximum number of hatchery-reared steelhead trout while releasing wild Middle Fork fish.

Future management considerations for the Middle Fork Salmon River steelhead trout population are discussed.

Author:

Russ Thurow
Senior Fishery Research Biologist

INTRODUCTION

Historically large runs of wild steelhead trout (Salmo gairdneri) returned annually to Idaho's abundant rivers and streams. The construction of dams has eliminated nearly 50% of the steelhead trout's original habitat by totally blocking runs in the Boise, North Fork Clearwater, Payette, Upper Snake and Weiser rivers.

The Middle Fork Salmon River is one of three major Idaho rivers which sustain wild steelhead trout (steelhead) runs unaltered by hatchery propagation. Historically, the Middle Fork supported substantial runs of steelhead which probably exceeded an annual spawner escapement of 10,000 fish. During the last decade, hydroelectric dams on the Columbia and Snake rivers have severely reduced survival of migrating steelhead, and Middle Fork spawning escapements have diminished from approximately 5,000 in 1970-71 to 500 (or less) in 1975-76 (Jeppson and Ball 1979). Consequently, the Middle Fork has been closed to steelhead fishing since 1974 in an attempt to sustain the wild steelhead stock.

Although several biologists have evaluated the cutthroat trout (Salmo clarki) resources of the Middle Fork drainage (Mallet 1963, Ortman 1971, Ball and Jeppson 1980), very little data has been collected on its steelhead resources. Preliminary work on steelhead was conducted in 1980 (Reingold 1981). In 1981, the Idaho Department of Fish and Game initiated an intensive 3-year fishery investigation. This project was designed to evaluate the current status of wild steelhead and to provide information to assist future management of the steelhead resource of the Middle Fork. Data on cutthroat trout (cutthroat) and other species was also collected.

Results of the 1981 and 1982 field work have been reported in job performance reports (Thurow 1982a, 1983). This report includes 1983 field data and a synopsis of the research program from 1981 to 1983.

Since 1981, significant changes in steelhead fishing regulations have been enacted on the Salmon River. The major objective has been to differentially harvest hatchery stocks, while increasing escapements of wild stocks.

During the fall 1980-spring 1981 fishery, no special limits or gear restrictions were set, although anglers were encouraged to voluntarily release wild steelhead. In the fall 1981, the possession limit was increased from two to four fish above the Middle Fork, and in the spring 1982, the bag, possession and season limits were set at one, one, three, respectively, below the Middle Fork and two, four, six, respectively, above the Middle Fork. Barbless hooks were also required below the Middle Fork. In the fall 1982, a regulation was initiated in Section 4 (South Fork to Middle Fork) and a portion of Section 3 (Vinegar Creek to South Fork) which required anglers to use barbless hooks and release most wild steelhead based on a dorsal fin measurement (Thurow 1983). Steelhead larger than 94 cm were exempted to allow anglers to keep a "trophy" fish. In the spring 1983 these regulations were extended to include Sections 1 through 4. Most recently, regulations for the fall 1983-spring 1984

fishery were set to require release of wild fish, based on a dorsal fin height measurement, regardless of total length.

An ultimate goal of this program is to restore angling opportunities for wild steelhead within the Middle Fork drainage. The baseline studies have provided management alternatives and a means of monitoring the status of the population. Descriptions of fish populations in Middle Fork tributaries also provide management biologists with data for technical assistance.

DESCRIPTION OF STUDY AREA

The Middle Fork of the Salmon River flows through a remote area of central Idaho and for most of its length, lies within the Frank Church River of No Return Wilderness Area. From its origin at the confluence of Bear Valley and Marsh creeks, the Middle Fork flows north-northeast for 171 km through the Salmon River mountains and joins the Salmon River 92 km below Salmon, Idaho (Fig. 1).

The earliest known human inhabitants of the Middle Fork were likely Paleo Indians 10,000 years ago (Knudson et al. 1982). White men initially described the drainage in 1824 when Alexander Ross traveled along Marsh Creek. Carrey and Conley (1980) and Knudson provide detailed discussion of the prehistory and human history of the Middle Fork drainage.

Mallet (1963), Minshall et al. (1981) and Thurow (1982a) provide detailed descriptions of study area's topography, climate, vegetation, stream discharge, water quality and recreational use.

Peak stream discharges occur during a two- to six-week period in May and June as a result of snowmelt. Spring runoff extends over four months, with a base flow over the remaining eight months. Flows decrease throughout summer and increase with the onset of winter precipitation.

Mean annual discharge equals 43.2 m³/sec (1973-1980). Flows during the past five years varied considerably between years, ranging from a 1977 mean annual flow of 16.1 m³/sec, and a maximum discharge of 52.7 m³/sec, to a mean flow of 43.9 m³/sec, and a maximum discharge of 255.1 m³/sec in 1980 (USGS 1977-1980). The gauge at Middle Fork Lodge was discontinued in 1981.

Road access exists to Dagger Falls and at the Middle Fork's confluence with the Salmon River. Although headwaters of a few tributaries are accessible via unimproved roads, the lower 156 km of the Middle Fork is accessible by air, float craft, or trail only. Ortmann (1969) observed that the Middle Fork has attained national prominence as a recreational stream since it offers outdoor enthusiasts opportunities in whitewater boating, angling, hunting, or passive enjoyment of rugged scenery. In 1983, 7,943 people floated the Middle Fork (Challis National Forest 1984), compared to 625 in 1962 (Ortmann 1969).




Figure 1. Middle Fork Salmon River drainage, Idaho.

Although most of the Middle Fork drainage and its aquatic habitat lies in a pristine wilderness, human activity has significantly altered sections of several tributaries. Precious metal mining has caused extensive sediment transport to Monumental and Big creeks. In July 1981, activities at the Golden Reef Joint Venture Mine resulted in an influx of sediment pond wastewater to Mule and Monumental creeks. In October 1983, several tons of settling pond sludge spilled into Mule Creek, tributary to Monumental Creek. A fish habitat survey was conducted in Monumental Creek on October 19, 1983, by Idaho Department of Fish and Game and U.S. Forest Service personnel (Burns 1983). The biologists measured embeddedness above and below the confluence of Mule Creek and determined there is 50% less available fish habitat below Mule Creek than above as a result of man-caused sedimentation. The State Land Board filed suit against Golden Reef, and an administrative action was filed before the Board of Health and Welfare. Rehabilitation efforts and additional water quality monitoring were ordered by the Board of Health and Welfare.

Extensive placer mining has occurred along the upper reaches of Loon Creek. In 1983, the Loon Creek Mining Company submitted additional proposals to activate a placer mining operation adjacent to Loon Creek near the Oro Grande townsite. The proposed mining operation has the potential to damage water quality and fish habitat in Loon Creek (Thurrow 1982b).

On Camas Creek, livestock use has degraded riparian habitats in Meyers Cove. The Cobalt District of the Salmon National Forest has identified the need for additional control of livestock use in Meyers Cove, and fencing programs have been initiated as part of a new grazing management system. Three placer mine operating plans have been filed for operations on Silver Creek (tributary to Camas Creek).

Grazing by livestock has degraded riparian and instream habitat in Marsh, Bear Valley and Elk creeks (Thurrow 1983). Past gold dredging in Bear Valley Creek has deposited sediment and eliminated fish habitats.

Within the Marble Creek drainage, Coeur d'Alene Mines Corporation has filed a Notice of Intent with the Payette National Forest to operate a 10- to 20-year mining project. The proposal would include digging ore from an open pit mine and operating a cyanide process gold extraction mill.

In 1980, the man-caused Mortar Creek fire burned over 26,000 hectares and extended along 40 km of the Middle Fork, including several tributaries. Minshall et al. (1981) monitored the effects of the fire and found that Little Loon Creek sustained the greatest damage. Heavy runoff and mass wasting transported massive amounts of sediment, scoured the stream bed and caused large amounts of material to enter the main river.

OBJECTIVES

To document the principal steelhead trout spawning areas in the Middle Fork Salmon River drainage and to assess adult escapement.

To collect biological data characterizing the size, sex ratio and origin (wild or hatchery) of steelhead trout spawners.

To assess the abundance, distribution and population structure of cutthroat trout and juvenile steelhead trout in the Middle Fork Salmon River and tributaries.

To genetically characterize fish from Middle Fork Salmon River tributaries in order to compare them to each other and to other Idaho steelhead stocks.

To evaluate the timing and movement of wild steelhead trout in the main Salmon River and in the Middle Fork Salmon River and tributaries.

To assess the contribution of wild Middle Fork Salmon River steelhead trout to the main Salmon River sport fishery downstream from the Middle Fork.

RECOMMENDATIONS

1. Wild steelhead trout stocks are unique in Idaho. The Middle Fork Salmon River is one of three drainages which sustain steelhead trout unaltered by hatchery-reared stocks. Continued management for the production and preservation of the indigenous stock is the recommended alternative for restoration of steelhead trout sport fishing opportunities to the Middle Fork Salmon River.
2. Differential harvest regulations initiated in 1982 have been successful in increasing escapements of wild steelhead trout. Maintenance of these regulations on the Salmon and Snake rivers will aid restoration of the Middle Fork Salmon River steelhead trout population.
3. Optimal escapement of wild steelhead trout, defined as the number of spawners required to fully seed the available habitat with parr, is recommended for the Middle Fork Salmon River. Calculations based on the application of steelhead production data to available habitat suggest a spawning escapement goal of approximately 8,000, with a range from 6,000 to 11,500 depending on spawning success.
4. Escapement estimates for wild steelhead trout are critical for future management decisions. The angler creel census initiated on the Salmon and Snake rivers in 1984 will quantify escapements of wild steelhead trout to the Middle Fork Salmon River using wild fish: hatchery fish ratios and hatchery escapements. Estimates can be compared to escapement goals to monitor the population status. Corroborative field data can be collected with a small expenditure of effort. Snorkeling counts of selected tributary transects and redd counts of index areas should be conducted annually. Selection of tributary transects can be coordinated with research personnel. Recommended procedures for and location of redd index surveys are in Appendix A.

5. Maintenance of fisheries habitats in a pristine condition will assist the restoration of wild steelhead trout. An aggressive stance for habitat protection is warranted when reviewing proposals which have the potential to degrade aquatic habitats. Fish populations will benefit if corrective measures are applied to restore aquatic habitats which have been degraded. Impacted areas listed in the Description of Study Area require restoration efforts.
6. Section 9(a) of the Central Idaho Wilderness Act addresses a ban on dredge and placer mining. The wording and intent of the Act are subject to interpretation. A legal opinion is needed to resolve the intent of the Act.
7. Accessible sections of Indian, Loon and Pistol creeks appear to receive considerable angling effort. A periodic creel census would define the angler effort and harvest.
8. With the exception of Big Creek, anglers can harvest fish of any size in tributaries. Steelhead trout parr apparently comprise a majority of the fish caught in tributaries. A 200 mm minimum size limit would restrict the harvest of most steelhead trout parr. The regulation would also enable anglers to harvest residualized steelhead trout, resident rainbow trout and larger cutthroat and bull trout.

METHODS

Spawning Area Surveys

Commencing in March and extending to the third week in May, we visually surveyed and fished sections of tributaries to the Middle Fork during 1981, 1982 and 1983. We surveyed 262 km of stream in 1981, 106 km in 1982 and 337 km in 1983. Thirteen streams were surveyed by both ground and aerial methods. We mapped stream sections, counted steelhead spawners and redds, observed spawning behavior and recorded biological data on fish captured or observed. Since most steelhead apparently remain on redds from one to three days and migrate from spawning areas soon after spawning (T. Johnson, Washington Department of Game, pers. comm.; Reingold 1964), it is unlikely we counted any fish more than once.

Springtime visual surveys of spawning areas do not provide a reliable estimate of actual spawner abundance. Water conditions are subject to change, and even during excellent conditions, only a portion of the spawners in a stream reach are visible. However, surveys do provide important information on the timing and location of steelhead spawning in addition to biological data characterizing spawners.

Survey conditions were excellent in 1981 and 1983 and poor in 1982.

Juvenile Distribution and Abundance

We used snorkeling counts of fish in previously established transects to assess the abundance of juvenile steelhead. Counts were made on cloudless days between 0930 and 1630 when visibility was maximum. Several researchers (Northcote and Wilkie 1963, Goldstein 1978, Griffith 1980) have concluded that reliable estimates of fish abundance can be obtained by underwater counts. We completed our counts in July and August because juvenile steelhead maintain specific daytime stations and home ranges in summer (Edmundson et al. 1968). Underwater census techniques were ideal for surveying the streams in the Middle Fork drainage. Everest (1969) quantitatively described habitat selected by juvenile steelhead. We reviewed his descriptions and selected transects exhibiting abundant rubble-boulder substrates, moderate or faster velocities and run-slick qualities (Thurow 1982a). Pools, riffles and shallow runs were not selected because they are not preferred habitats. Most transects contained abundant rubble-boulder pocket water habitat.

Juvenile steelhead were classified by length at Age I (70-130 mm), Age II (130-200 mm) and Age III (>200 mm) using a classification similar to Everest (1969). I did not attempt to count young-of-the-year salmonids (<70 mm) since they were indistinguishable by species and timing of complete emergence was unknown. It is likely that most fish larger than 250 mm were residualized steelhead or resident rainbow trout. Idaho Cooperative Fishery Unit personnel measured lengths of 1,592 wild steelhead smolts at Lower Granite Dam in 1977 (unpublished). Smolts ranged from 120 to 290 mm and most (89% to 96% for two groups) ranged from 170 to 250 mm. One to six percent exceeded 250 mm and only 1% of the total 1,592 exceeded 270 mm. Since we sampled Middle Fork fish in August, they would have continued to grow until late October, with additional growth in spring prior to smoltification. Everest (1969) estimated a length increase of 9 mm per month in the third summer of a steelhead's freshwater rearing state. Consequently, a juvenile steelhead 250 mm long in mid-August would exceed 270 mm prior to the following spring, and it is unlikely a fish of that size would undergo smoltification and migrate. Within this report, I assumed that below barriers all rainbow trout or steelhead parr less than 250 mm were juvenile steelhead trout. It was further assumed that all rainbow trout or steelhead parr larger than 250 mm were non-smolting residualized steelhead or resident rainbow trout. Rainbow trout observed above barriers were classified as resident rainbow trout. A resident rainbow trout population often remains when steelhead are blocked from an area (Simpson and Wallace 1982).

Total numbers of other game fish were counted by species. Within the Middle Fork, the number of cutthroat larger than 300 mm was recorded. Within tributaries, cutthroat were recorded in 100 mm size groups. The presence of mountain whitefish (Prosopium williamsoni) and nongame species was noted.

To assess length frequency distributions of steelhead and other game fish, we used barbless flies and lures to capture fish in the Middle Fork and tributaries. Species, total length, date and location of capture were recorded for all fish caught.

Middle Fork Salmon River

Twenty longitudinal transects had been established in 1980 at sites we considered good steelhead habitat (rubble-boulder pocket water) between Boundary Creek and the Salmon River (Reingold 1981). Chapman and Bjornn (1969) used the term "rubbly-glide" to describe such habitat. Using a wetsuit and snorkel, I floated two separate glides (visible corridors) down each transect and enumerated all fish by species (Thurow 1982a). One glide was made close to the shoreline and the second near midstream. A second diver made consecutive passes down each glide approximately five minutes later and the maximum count was used. Following the counts, we measured the total length of each transect and recorded water temperatures. Visibility was determined by measuring the distance a diver was able to see a brass scale underwater. The area snorkeled was calculated by the formula:

$$\text{Surface area} = (2V) (L) (G)$$

Where: V = Visibility
L = Total transect length
G = Number of glides snorkeled.

Each transect was photographed and physical descriptions and channel characteristics recorded.

Tributaries

We established and surveyed transects in 12 tributaries. Streams were separated into upper, middle and lower sections. Within each section, we established five transects of similar length at sites we considered good steelhead habitat. Using a wetsuit and snorkel, I proceeded upstream through each transect and counted all fish by species (Thurow 1982a).

After completing the counts, we measured the physical dimensions of each transect including total length and depth and width at 10 m intervals. We photographed each transect and recorded water temperatures, substrate, channel characteristics and riparian vegetation.

Origin of Middle Fork Steelhead

We used angling and electrofishing gear to collect juvenile steelhead in Big, Loon and Marble creeks in 1981 and 1982. Samples were packed in

dry ice and frozen prior to analysis. Laboratory personnel extracted a piece of skeletal muscle, the liver and eye fluid for electrophoretic analysis. Samples were screened for several genetic loci (Thurow 1982a) and 24 to 30 were consistently resolved.

Otolith nuclei were investigated as a means of distinguishing juvenile steelhead from juvenile rainbow trout in Big and Loon creeks using techniques described by Thurow (1982a).

Adult Steelhead Movements

During the spring and fall, we captured and tagged adult steelhead in the Middle Fork and main Salmon rivers. We used barbless lures, flies and bait to capture fish, then attached numbered metal tags to their mandibles. Total length, dorsal fin height, sex, tag number and date and location of capture were recorded before releasing the fish.

To obtain timing and movement data, we refished areas of the Middle Fork and Salmon rivers. We also checked for tags on all angler-creeled fish we encountered on the main Salmon River. Informational signs explaining the tag program were posted along the main Salmon River, and tag deposit boxes were placed locally. A brief history of the tagged fish was sent to each angler who returned tag recovery information.

We made several float trips in the fall to document the location of adult steelhead downstream from the Flying "B" Ranch. Angling and snorkeling gear were used to locate fish.

Main Salmon River Sport Fishery

We annually operated a check station on the Salmon River at the mouth of the Middle Fork to monitor the steelhead fishery below that point. Part of the season was divided into three two-week intervals as follows: (1) October 10-23, (2) October 24 to November 6, and (3) November 7-20. Within each interval, we randomly selected three weekdays and two weekend days. Holidays were also censused.

On each census day, a clerk interviewed all anglers leaving by the single access road from 1000 hours to darkness. The clerk recorded numbers of anglers, stream section fished and numbers of fish creeled and released. Total length, dorsal fin height, sex and origin (wild or hatchery) were recorded.

We estimated the total harvest of steelhead per interval as follows:

$$\text{Catch} = \bar{x}_1 \text{ WD} + \bar{x}_2 \text{ WE} + \bar{x}_3 \text{ H}$$

- \bar{x} = mean catch per weekday
 \bar{x}_2 = mean catch per weekend day
 \bar{x}_3 = mean catch per holiday
WD = number of weekdays per interval
WE = number of weekend days per interval
H = number of holidays per interval

A jet boat was used to monitor the steelhead fishery in the roadless area below Corn creek (11 km below the Middle Fork). The roadless area fishery also was monitored by interviewing anglers at the check station, conducting field checks of anglers and gathering data compiled by cooperating outfitters.

RESULTS

Biological Characteristics

Both anadromous (steelhead) and nonmigratory (resident) rainbow trout (Salmo gairdneri) are indigenous to the Middle Fork Salmon River. These rainbow trout may be analogous to the redband trout (Salmo sp.) described by Behnke (1979).

The fish fauna of the Middle Fork is represented by five families (Catostomidae, Cottidae, Cyprinidae, Petromyzontidae and Salmonidae), 10 genera and 16 species (Table 1). Only one species (brook trout, Salvelinus fontinalis) is nonindigenous and it is found only in isolated areas.

Macroinvertebrates are abundant in the Middle Fork and tributaries and are represented by five principal orders consisting of 81 taxa (Minshall et al. 1981). Generally, tributary streams supported 40+ and the mainstream 25+ taxa on a seasonal basis. The orders Ephemeroptera, Diptera and Trichoptera were the most prevalent organisms collected in both the mainstem and tributaries. Plecoptera were also prevalent in tributaries.

Spawner Characteristics and Densities

Seventy-eight steelhead spawners and 80 redds were observed for an average of one spawner or redd per 5 km surveyed (Appendix B). Spawning activity varied by stream and timing of surveys. Peak densities of spawners and redds were observed in Big, Camas and Loon creeks.

Table 1. Fishes present in the Middle Fork Salmon River drainage.

Common name	Scientific name	Middle Fork	Status by location	Tributaries
Rainbow-steelhead trout	<i>Salmo gairdneri</i>	abundant	abundant	abundant
Westslope cutthroat trout	<i>Salmo clarki lewisi</i>	abundant	abundant	common
Chinook salmon	<i>Oncorhynchus tshawytscha</i>	currently depressed	currently depressed	currently depressed
Bull trout	<i>Salvelinus confluentus</i>	common	common	common
Mountain whitefish	<i>Prosopium williamsoni</i>	abundant	abundant	common
Northern squawfish	<i>Ptychocheilus oregonensis</i>	common in lower 32 km	common in lower 32 km	uncommon
Redside shiner	<i>Richardsonius balteatus</i>	common in lower 32 km	common in lower 32 km	uncommon
Bridgelip sucker	<i>Catostomus columbianus</i>	unknown	unknown	unknown
Largescale sucker	<i>Catostomus macrochellus</i>	unknown	unknown	uncommon
Longnose dace	<i>Rhinichthys cataractae</i>	common	common	common
Speckled dace	<i>Rhinichthys osculus</i>	unknown	unknown	unknown
Shorthead sculpin	<i>Cottus confusus</i>	unknown	unknown	unknown
Mottled sculpin	<i>Cottus bairdi</i>	unknown	unknown	unknown
Torrent sculpin	<i>Cottus rhotheus</i>	unknown	unknown	unknown
Pacific lamprey	<i>Entosphenus tridentatus</i>	unknown	unknown	unknown
Brook trout (introduced)	<i>Salvelinus fontinalis</i>	absent	absent	headwaters of Marsh & Big creeks

Spawning activity commenced in early April, and peak activity occurred in May (Fig. 2). Conditions were too turbid for surveys after late May. We observed most of the spawners and redds in Monumental and Big creeks between May 17-20, 1983.

Steelhead spawners ranged from 61-91 cm, and sex ratios averaged 1:1.

Individual tributaries surveyed in 1983 are listed in the following section. Tributaries surveyed in previous years are discussed in Thurow (1982a, 1983). Additional data for previously surveyed streams are presented in Appendix B.

Brush Creek

Brush Creek enters the Middle Fork 47 km above the mouth and is 9.7 km long to the confluence of its north and south forks (Fig. 3). A hydropower diversion for the Flying "B" Ranch creates a barrier to fish passage approximately 1.6 km above the mouth. A natural barrier exists approximately 0.8 km above the diversion dam. Suitable spawning substrate exists both below and above the diversion dam and barrier. No spawning activity was observed.

Marble Creek

Marble Creek enters the Middle Fork 101 km above the mouth and is 39 km long (Gebhards 1959). Average discharge is approximately 3.5 m³/sec. Survey conditions were unfavorable in 1983, and we observed one spawner and redd (Fig. 4). Extensive spawning area is present throughout the drainage, including several smaller tributaries.

Monumental Creek

Monumental Creek enters Big Creek approximately 50 km above the mouth and is Big Creek's largest tributary. Lower sections of Monumental Creek were surveyed in 1981 (Thurow 1982a). In 1983, we surveyed the entire drainage from Roosevelt Lake to Big Creek and observed large densities of spawners and redds (Fig. 5). Monumental Creek supported the largest numbers of steelhead spawners we observed in the Big Creek drainage. Densities of spawners and redds were the largest we observed in any Middle Fork tributary from 1981 to 1983 (one spawner or redd each 1.2 km surveyed).

Sheep Creek

Sheep Creek enters the Middle Fork 49 km above the mouth and is 8.1 km long to the confluence with its south fork (Fig. 3). A steep gradient

▲ CAMAS N: 46
 ■ BIG N: 73
 ● LOON N: 88




Figure 2. Percent of steelhead spawners and redds (combined) observed in Big, Camas and Loon creeks during specified periods, 1968-1983.



Figure 3. Spawning ground survey map of Brush and Sheep creeks, Middle Fork Salmon River, Idaho, 1983.




Figure 4. Spawning ground survey map of Marble Creek, Middle Fork Salmon River, Idaho, 1983.




Figure 5. Spawning ground survey map of Monumental Creek, Middle Fork Salmon River, Idaho, 1983.

section, approximately 6 km above the mouth, may create an impassible barrier to steelhead. Suitable spawning area is prevalent in the lower 3 km and scattered above that point. No spawning activity was observed.

Sulphur Creek

Sulphur Creek enters the Middle Fork 152 km above the mouth and is 31 km long (Gebhards 1959). Average discharge is approximately $1.7 \text{ m}^3/\text{sec}$. Sulphur Creek is a principal spawning and rearing area for chinook. Suitable steelhead trout spawning substrate is also present (Fig. 6). A barrier to fish migration exists 17.7 km above the mouth. No spawning activity was observed.

Wilson Creek

Wilson Creek enters the Middle Fork 37 km above the mouth and is 22.5 km long (Gebhards 1959). Flow is approximately $1.8 \text{ m}^3/\text{sec}$. A large rock slide and "blow out" occurred in the 1950's above Alpine Creek. The event created a barrier to fish movement 6 km above the mouth. Suitable spawning area is present throughout the drainage and some excellent substrate exists in upper reaches of the stream (Fig. 7). No spawning activity was observed.

Yellowjacket Creek

Yellowjacket Creek enters Camas Creek 8 km above the mouth and is 40 km long (Gebhards 1959). Average discharge is approximately $1.7 \text{ m}^3/\text{sec}$. The stream contains suitable spawning substrate above the Yellowjacket Mine for several miles (Fig. 8). A constricted area of the channel, 2.4 km above Camas Creek, was believed to be a passage barrier, and in 1976, the USDA-FS removed a barrier at the site. No spawning activity was observed.

Juvenile Distribution and Abundance

Middle Fork

Numbers and lineal densities (fish/100 m) of juvenile steelhead in the Middle Fork transects increased nearly three-fold from 1980 to 1981 and were similar from 1981 to 1983 (Table 2, Fig. 9). In contrast, densities of juvenile steelhead per unit surface area (fish/100 m^2) were similar in 1981 and 1982 and nearly doubled in 1983. A two-factor analysis of variance was used to test for differences in densities between years. For lineal densities (fish/100 m) significantly ($p < 0.05$) more steelhead were

Table 2. Numbers and densities of juvenile steelhead observed in Middle Fork Salmon River transects, 1983.

Transect	1983 count		Total 1983	Steelhead per 100 m		Steelhead per 100 m ² 1983
	Age I+	Age II+		Age III+	1983	
1	4	3	8	6.2	.49	
2	3	3	7	6.7	.53	
3	3	7	12	6.6	.47	
4	1	0	2	2.2	.16	
5	9	13	26	10.5	.75	
6	3	8	12	5.2	.66	
7	3	5	11	3.0	.39	
8	3	19	24	11.4	1.47	
9	5	11	19	7.0	.86	
10	2	5	7	6.7	1.19	
11	4	11	17	5.3	.80	
12	2	3	5	2.0	.31	
13	2	1	4	1.6	.28	
14	1	10	14	6.1	.66	
15	3	3	11	5.2	.57	
16	1	3	5	2.2	.24	
17	1	6	10	5.0	.54	
18	2	3	6	1.8	.22	
19	3	5	12	3.7	.38	
20	--	2	4	2.4	.25	
Totals	55	121	216	$\bar{x} = 4.9$	$\bar{x} = .540$	




Figure 6. Spawning ground survey map of Sulphur Creek, Middle Fork Salmon River, Idaho, 1983.




Figure 7. Spawning ground survey map of Wilson Creek, Middle Fork Salmon River, Idaho, 1983.




Figure 8. Spawning ground survey map of Yellowjacket Creek, Middle Fork Salmon River, Idaho, 1983.




Figure 9. Number and density of juvenile steelhead observed in Middle Fork Salmon River transects, 1980-1983.

observed from 1980 to 1983. Densities per unit surface area (fish/100 m²) were similarly tested and no significant (P<0.05) differences occurred in Age I+ steelhead densities from 1981 to 1983 although significant (P<0.05) differences occurred in Age II+ and total steelhead densities during the same period.

Although lengths of individual transects remained similar from 1980-1983, visibility varied significantly (Appendix C). Transects ranged from 65 to 252 m and visibility ranged from 5.4 to 7.5 m in 1981 and 7.8 to 9 m in 1982. In August 1983, severe rains created turbid water conditions in several tributaries (particularly Little Loon and Marble creeks), and visibility decreased to 2.8 to 4.8 m below Indian Creek. Water temperature from Boundary Creek to the mouth varied from 12 to 19 C in 1981, 12 to 17 C in 1982 and 13 to 19.5 C in 1983.

Angling proved to be an effective technique for collecting fish species composition and length frequency data. We caught and released 1,942 game fish in the Middle Fork during July and August 1980-1983 (Table 3). In 1982, 760 game fish were caught in 24 man days of fishing. Artificial flies were consistently the most effective terminal tackle.

In 1983, juvenile steelhead comprised 45% of the catch, cutthroat 51% and rainbow x cutthroat trout hybrids, bull trout (Salvelinus confluentus) and chinook salmon (Oncorhynchus tshawytscha) less than 1% (Table 3). In comparison, juvenile steelhead comprised 44% of the fish observed by snorkeling, cutthroat 33%, chinook salmon 21% and bull trout 2%. Transects were selected in optimal steelhead habitat while hook-and-line sampling covered all habitats. The small size of juvenile chinook salmon and the sedentary behavior of whitefish resulted in both species not being caught in proportion to their abundance.

The age-frequency of juvenile steelhead observed by snorkeling varied among years (Table 4). Age II+ and I+ fish predominated with smaller percentages of Age III+ fish. Hook-and-line gear was ineffective in sampling steelhead less than 130 mm (Age I+), but was effective in sampling larger fish. Length frequencies of steelhead caught by angling were similar from 1980 to 1983 (Appendix D). We measured 893 steelhead which ranged from 90 to 370 mm (Fig. 10).

We observed cutthroat trout in 57 of 60 transects sampled. Cutthroat were most abundant in transects between Rapid River and Tappan Falls and least abundant below Big Creek. Numbers of cutthroat were similar from 1981 to 1983 and averaged 159 per 20 transects (Table 5). Densities of cutthroat ranged from 2.4 to 3.7 fish/100 m and averaged 3.1 fish/100 m. Cutthroat per unit surface area increased from 0.19 to 0.35 fish/100 m² from 1981 to 1983. Cutthroat abundance is affected by seasonal movements as Mallet (1963) observed.

Length frequencies of hook-and-line caught cutthroat were similar from 1980 to 1983 (Appendix D). We measured 1,009 cutthroat which ranged from 130 to 430 mm (Fig. 10). Forty-eight percent of the fish exceeded 300 mm and 0.2% exceeded 400 mm. We observed a decline in the abundance of cutthroat larger than 300 mm from 1980 to 1983 in our hook-and-line samples. The percentage of large cutthroat (>300 mm) also declined in our snorkeling surveys from 62% in 1980 to 53% in 1983 (Table 5).

Table 3. Number of fish sampled by hook-and-line and snorkeling in the Middle Fork Salmon River, July-August 1980-1983.

Year	Steelhead	Cutthroat	Hook-and-Line			Chinook salmon	Mountain whitefish	Sub-totals
			Hybrid (rainbow x cutthroat)	Bull trout				
1980	167	190	8	6	1	3	375	
1981	126	133	6	0	1	2	268	
1982	311	396	7	4	0	42	760	
1983	266	265	5	1	1	1	539	
Total	870	984	26	11	3	48	1,942	
Percent	45	51	1	<1	<1	2		

Snorkeling								
1981	200	143	a	10	18	b	371	
1982	215	194	a	5	161	b	575	
1983	216	139	a	11	115	b	481	
Total	631	476	a	26	294	b	1,427	
Percent	44	33	--	2	21	--		

a - Not identified.
b - Not enumerated.

Table 4. Age frequency of juvenile steelhead observed in twenty Middle Fork Salmon River transects, 1980-1983.

=====

Year	Percent of juvenile steelhead observed		
	Age I+	Age II+	Age III+
1980	36	53	11
1981	45	42	13
1982	45	43	12
1983	25	56	19

=====

Table 5. Numbers of fish (non-steelhead) observed in Middle Fork Salmon River transects, August 1983.

Section	Transect	Cutthroat		Juvenile chinook salmon		Bull trout	Mountain whitefish	Suckers	Nongame fish (+,-)	Redside shiner
		Total	No. >300 mm	Total	No. >300 mm	trout	whitefish	Suckers	Northern squawfish	
I	1	6	6	0	0	0	+	-	-	-
	2	6	4	13	0	0	+	-	-	-
	3	8	4	2	0	0	+	-	-	-
	4	2	2	2	2	2	+	-	-	-
	5	28	14	12	4	4	+	-	-	-
	6	14	9	3	0	0	+	-	-	-
II	7	7	4	12	0	0	+	-	-	-
	8	12	8	1	1	1	+	-	-	-
	9	9	2	1	1	1	+	-	+	-
	10	0	0	1	0	0	+	-	-	-
	11	4	2	11	0	0	+	-	-	-
	12	3	2	7	0	0	+	+	-	-
III	13	0	0	11	0	0	+	+	-	-
	14	10	5	9	0	0	+	+	-	-
	15	4	1	2	0	0	+	+	+	-
	16	8	3	0	0	0	+	+	+	+
IV	17	0	0	18	0	0	+	+	+	-
	18	2	1	8	1	1	+	+	+	-
	19	7	2	2	0	0	+	+	+	-
	20	9	4	0	2	2	+	+	+	-
Total numbers		139	73	115	11					
Number of transects where species present:		17	17	17	6	20	9	7	1	

In addition to steelhead and cutthroat, we also observed and captured other game fish and nongame species. Although we did not record their abundance, mountain whitefish were the most abundant game fish as Corley (1972) and Jeppson and Ball (1979) also observed. Mountain whitefish were present in all 80 snorkeled transects from 1980 to 1983. We captured 48 mountain whitefish with angling gear and 94% exceeded 280 mm.

Bull trout were often difficult to observe, and snorkeling did not provide a reliable estimate of their abundance. We counted 26 bull trout in 60 transects (Thurow 1982a, 1983) (Table 5). Corley (1972) and Jeppson and Ball (1979) observed only 14 and 1 bull trout, respectively, in the 21 transects they snorkeled in 1971 and 1978. We also caught few bull trout with artificial flies, comprising less than 1% of the 1,942 fish caught (Table 3). In 1959 and 1960, the percentage of bull trout in the catch ranged from 4 to 14% (Mallet 1963). The fish we caught in summer ranged from 150 to 360 mm. We also captured bull trout during spring and fall surveys ranging up to 560 mm. As Jeppson and Ball (1979) observed, most bull trout in the Middle Fork are caught during the spring and fall months. Mallet (1963) reported that bull trout migrate into the Middle Fork from the mainstem Salmon River during the fall and winter.

Juvenile chinook salmon were in low abundance and we observed 294 in 60 transects (Table 5). Captured chinook salmon ranged from 75 to 115 mm.

We could not differentiate rainbow x cutthroat trout hybrids while snorkeling. We captured 28 with hook-and-line, ranging from 180 to 410 mm. Forty-three percent exceeded 300 mm and 11% exceeded 350 mm.

Northern squawfish (Ptychocheilus oregonensis) and suckers (Catostomus sp.) were most abundant in river sections below Tappan Falls (Table 5). We observed northern squawfish in only 4 of 33 transects snorkeled above the falls. Redside shiners (Richardsonius balteatus) were observed in one transect each of the three years.

Tributaries

Juvenile steelhead were much more abundant in tributaries than in the Middle Fork. Tributaries provide the principal rearing habitat for steelhead in the drainage. We snorkeled 153 transects in twelve major tributaries and observed 2,263 juvenile steelhead for an average of nearly 15 per transect (Table 6).

Most transects ranged from 30 to 50 m long and varied considerably in width depending on the streams (Appendix E). Rubble was the predominant substrate, followed by gravel and boulders. Sand and silt comprised a small proportion of the substrate in most stream sections. Bear Valley Creek, upper Marsh Creek and Elk Creek contained the largest percentage of sand substrate. Riparian vegetation consisted of grasses, sedges, various brush and shrubs. Forest canopies of pine and fir were generally sparse. Water temperatures ranged from 9 to 17 C in individual tributary sections.

Table 6. Numbers and sizes of fish observed by snorkeling in Middle Fork Salmon River tributaries, July-September, 1983.

Stream	Juvenile Steelhead				Cutthroat (mm)				Total	Total >300	Juvenile chinook salmon	Mountain whitefish	Bull trout	Age 0 salmonids	Steelhead per 100 m ²
	Age I+	Age II+	Age III+	Total	100	100-200	200-300	>300							
Marble Creek															
Upper	2	7	3	12	25	54	25	0	104	0	0	0	0	+	1.7
Sunnyside	0	0	0	0	1	2	2	0	5	0	0	0	0	+	0
Cornish	0	0	0	0	1	2	5	0	8	0	0	0	0	-	0
Totals	2	7	3	12	27	58	32	0	117	0	0	0	0	+	
Sheep Creek															
Lower	34	43	13	90	0	10	7	0	17	0	0	0	9	-	10.5
Sulphur Creek															
Lower	24	20	6	50	0	3	1	0	4	17	0	0	0	+	2.2
Upper	0	0	0	0	3	10	8	0	21	0	0	0	0	±	0
Totals	24	20	6	50	3	13	9	0	25	17	0	0	0	+	
Wilson Creek															
Lower	44	61	9	114	0	0	0	0	0	0	0	0	0	-	7.8
Upper	78	90	19	177	0	0	0	0	0	0	0	0	2	-	13.6
Totals	122	141	28	281	0	0	0	0	0	0	0	0	2	-	
Yellowjacket Creek															
Lower	65	76	16	157	0	0	0	0	0	0	0	0	0	-	9.9
GRAND TOTALS	247	287	66	600	30	81	48	0	159	17	0	0	11	+	

-Absent
+Present




Figure 10. Length frequency of juvenile steelhead and cutthroat trout caught by angling in the Middle Fork Salmon River, 1980-1983.

Sections of Big, Camas, Loon and Wilson creeks supported the largest numbers of juvenile steelhead per transect. Based on angling surveys, lower sections of Marble Creek contain similar, large numbers of juvenile steelhead, but turbid water conditions prevented a snorkeling survey of the stream in 1983. Similarly, turbid water conditions prevented us from adequately surveying juvenile steelhead densities in Monumental Creek.

All 12 of the major drainages we surveyed (Bear Valley, Big, Camas, Indian, Loon, Marble, Marsh, Pistol, Rapid River, Sheep, Sulphur and Wilson creeks) contained populations of juvenile steelhead. In lower Brush Creek, we also observed juvenile steelhead, but did not snorkel any transects.

We located barriers to anadromous fish migration on upper sections of Indian, Sheep, Sulphur and Wilson creeks and on the lower kilometer of Brush Creek. Both Wilson and Brush creeks supported populations of residualized steelhead (resident rainbow trout) above the barriers.

Densities of juvenile steelhead within accessible tributary sections ranged from 0.2 to 10.5 fish per 100 m² snorkeled (Table 6). Sections of Big, Camas, Loon, Pistol, Sheep and Wilson creeks supported the largest densities of juvenile steelhead.

Angling was also effective in collecting data on tributary fish populations. Juvenile steelhead comprised a majority of the fish caught, followed by cutthroat, bull trout and rainbow x cutthroat trout hybrids (Table 7). I did not include mountain whitefish and juvenile chinook salmon in calculations because angling was ineffective in sampling. We captured steelhead, cutthroat and bull trout in proportion to their abundance as reflected by similar ratios for angling and snorkeling samples. Of 4,275 fish sampled in tributaries, 81% were steelhead, 16% cutthroat and 3% bull trout.

Age frequencies of steelhead observed by snorkeling averaged 42% Age I+, 52% Age II+ and 6% Age III+ (Table 6). Age III+ steelhead were less abundant in tributaries than in the Middle Fork, where they comprised 11 to 19% of the steelhead annually.

Angling continued to be ineffective in capturing steelhead less than 130 mm. We measured 1,992 juvenile steelhead which ranged from 80 to 370 mm (Fig. 11). Fish larger than 250 mm were probably resident rainbow trout or residualized steelhead. Fish of this size comprised 3% of the catch in tributaries compared to 9% in the Middle Fork.

Cutthroat were uncommon in most areas of the tributaries. We observed 415 cutthroat in 153 transects snorkeled for a mean of 2.7 per transect (Table 6). Fifty-six percent of the areas we snorkeled contained less than one cutthroat per transect, and 32% of the areas did not support any cutthroat. The upper section of Marble Creek supported the largest density of cutthroat trout (20.8/transect), followed by Indian (7.6/transect) and Pistol creeks (6.2/transect) (Table 8). Sixty-eight percent of the cutthroat we observed were sampled in the Indian, Marble

Table 7. Number of fish sampled by hook-and-line and snorkeling in Middle Fork Salmon River tributaries, July-August 1981-1983.

Year	Steelhead	Cutthroat	Bull trout	Hook-and-Line		Chinook salmon	Mountain whitefish	Totals
				Hybrid (rainbow x cutthroat)	Hook-and-Line			
1981	841	132	19	3	12	2	1,009	
1982	426	130	45	4	13	5	623	
1983	745	180	16	5	8	7	961	
Total	2,012	442	80	12	14	33	2,593	
Percent ^a	79	17	3	<1	0	0		

Snorkelling								
1981	1,079	68	11	b	92	c	1,250	
1982	584	188	62	b	302	c	1,136	
1983	600	152	11	b	17	c	787	
Total	2,263	415	84	b	411	c	3,173	
Percent ^a	82	15	3	--	--	--		

Grand								
Totals	4,275	857	164					
Percent ^a	81	16	3					

a - Chinook salmon and mountain whitefish were not included in calculations.
 b - Not identified.
 c - Not enumerated.

Table 8. Fish observed by snorkeling in Middle Fork Salmon River tributaries, July-August 1983.

Stream	Fish observed							Steelhead per 100 m ²
	Transect	Steelhead	Cutthroat	Bull trout	Juvenile chinook salmon	Surface area (m ²)	Steelhead per 100 m ²	
Camas Creek	M1	53	0	0	25	434	12.2	
	M2	18	0	1	4	502	3.6	
	M3	18	2	0	12	394	4.6	
	M4	27	0	0	2	373	7.3	
	M5	39	0	0	22	372	10.5	
Totals	155	2	1	65	2,075	$\bar{x}=7.5$		
Yellowjacket Creek (tributary to Camas Creek)	L1	23	0	0	0	257	9.0	
	L2	23	0	0	0	349	6.6	
	L3	26	0	0	0	268	9.7	
	L4	34	0	0	0	351	9.7	
	L5	51	0	0	0	358	14.3	
Totals	157	0	0	0	1,583	$\bar{x}=9.9$		
Marble Creek	U1	0	36	0	0	109	0	
	U2	0	20	0	0	112	0	
	U3	3	19	0	0	225	1.3	
	U4	3	17	0	0	134	2.2	
	U5	6	12	0	0	130	4.6	
Totals	12	104	0	0	710	$\bar{x}=1.7$		
Sunnyside Cr. Cornish Cr.		0	5	0	0	22	0	
		0	8	0	0	44	0	
Sheep Creek	L1	25	1	3	0	128	19.5	
	L2	23	1	4	0	206	11.2	
	L3	25	5	1	0	181	13.8	
	L4	6	5	0	0	167	3.6	
	L5	11	5	1	0	173	6.4	
Totals	90	17	9	0	855	$\bar{x}=10.5$		

Table 8. Continued.

Stream	Fish observed							Steelhead per 100 m ²
	Transect	Steelhead	Cutthroat	Bull trout	Juvenile chinook salmon	Surface area (m ²)	Steelhead per 100 m ²	
<u>Sulphur Creek</u>	L1	22	4	0	1	463	4.8	
	L2	23	0	0	7	470	4.9	
	L3	1	0	0	7	449	0.2	
	L4	4	0	0	0	404	1.0	
	L5	0	0	0	2	482	0.0	
Totals	50	4	0	17	2,268	$\bar{x}=2.2$		
Totals	U1	0	3	0	0	288	0	
	U2	0	5	0	0	204	0	
	U3	0	4	0	0	126	0	
	U4	0	6	0	0	212	0	
	U5	0	3	0	0	118	0	
Totals	0	21	0	0	948	$\bar{x}=0$		
<u>Wilson Creek</u>	L1	27	0	0	0	270	10.0	
	L2	21	0	0	0	209	10.1	
	L3	26	0	0	0	344	7.6	
	L4	21	0	0	0	328	6.4	
	L5	19	0	0	0	301	6.2	
Totals	114	0	0	0	1,452	$\bar{x}=7.8$		
Totals	U1	38	0	0	0	249	15.3	
	U2	29	0	2	0	330	8.8	
	U3	37	0	0	0	274	13.5	
	U4	42	0	0	0	251	16.7	
	U5	31	0	0	0	194	16.0	
Totals	177	0	2	0	1,298	$\bar{x}=13.6$		




Figure 11. Length frequency of juvenile steelhead and cutthroat trout caught by angling in Middle Fork Salmon River tributaries, 1981-1983.

and Pistol creek drainages. Areas of Big, Sheep and Sulphur creeks also contained abundant cutthroat trout populations. Headwater sections of Brush, Indian and Sulphur creeks support resident populations of cutthroat above barrier falls.

Densities of cutthroat within tributary sections ranged from 0.05 to 14.7 fish per 100 m² snorkeled (Table 8). Upper Marble Creek supported the largest cutthroat densities followed by sections of Indian, Pistol, Sheep and Sulphur creeks, which contained 1.3 to 2.2 fish per 100 m².

Length frequencies of cutthroat observed by snorkeling in tributaries averaged 10% for fish less than 100 mm, 47% 100 to 200 mm, 34% 200 to 300 mm and 9% larger than 300 mm (Table 6). Sections of Marble, Indian, Pistol and Sulphur creeks contained the largest number of fish less than 200 mm. Sections of Big, Loon and Pistol creeks contained the largest number of cutthroat which exceeded 300 mm. In comparison, resident cutthroat in upper Indian Creek averaged 4% less than 100 mm, 66% 100 to 200 mm, 30% 200 to 300 mm, and no cutthroat exceeded 300 mm.

Cutthroat captured with angling gear ranged from 60 to 410 mm and a majority (59%) ranged from 100 to 200 mm (Fig. 11). In comparison, resident cutthroat in upper Indian Creek rarely exceeded 250 mm (Thurrow 1983). Mature, resident cutthroat less than 200 mm were also captured in that section.

Bull trout were usually sympatric with cutthroat and were most prevalent in Indian, Loon, Pistol and Sheep creeks (Table 6). Bull trout were uncommon in most other tributary sections. We observed 84 in 153 transects. Fifty percent of the areas we snorkeled did not support any bull trout. We observed bull trout above barrier falls in Indian and Wilson creeks.

Eighty bull trout, ranging from 130 to 425 mm, were captured in tributaries (Fig. 12). A majority (80%) were less than 250 mm.

Mountain whitefish were present in all tributaries except Sheep Creek, and 68% of the areas we snorkeled contained mountain whitefish. Mountain whitefish were most prevalent in the deepest stream sections, where they exhibited schooling behavior. Young-of-the-year mountain whitefish were observed in Bear Valley and Marsh creeks. We captured 33 mountain whitefish ranging from 270 to 420 mm in the tributaries. Mountain whitefish were not captured in proportion to their abundance.

Juvenile chinook salmon were not common and we observed 411 in 153 transects (Table 6). Sixty-two percent of the areas contained juvenile chinook salmon. We observed juvenile chinook salmon in all tributaries except Sheep and Wilson creeks. We captured 14 juvenile chinook salmon with angling gear ranging from 80 to 140 mm. One precocial male captured in August measured 165 mm.

The only non-indigenous species we observed were brook trout (*Salvelinus fontinalis*), which had been introduced in Bear Valley, Big and Marsh creeks in the early 1900's. We observed 13 brook trout within five




Figure 12. Length frequency of bull trout caught by angling in Middle Fork Salmon River tributaries, 1981-1983.

transects in upper Big Creek, 5 brook trout, respectively, in single transects in Sack and Cache creeks (tributaries to Bear Valley Creek) and 167 brook trout in five transects in Marsh Creek above Capehorn Creek. A single, 45-meter-long transect in Marsh Creek near Kelly Creek contained 99 brook trout.

Other species, including dace (*Rhinichthys* sp.), sculpins (*Cottus* sp.) and suckers, were observed in tributaries while snorkeling, but we did not record their abundance.

Detailed descriptions of tributaries sampled in 1983 are provided in the next section. Data collected in tributaries sampled in 1981 and 1982 are provided in Thurow (1982a, 1983)

Bear Valley Creek

Newberry and Corley (1984) electrofished 13 tributaries to Bear Valley Creek and two sites within Bear Valley Creek in August 1982. Sampling sites were located in streams between Fir and Cassner creeks. Of the 15 sites surveyed, steelhead were collected in nine sites, bull trout in eight, brook trout in seven and chinook salmon in three. Three adult chinook salmon were sighted near the dredge mining area on August 16. Mountain whitefish were collected in six sites.

Steelhead were most abundant in Cache and Wyoming creeks and in an unnamed stream between Cold and Wyoming creeks. Only 51 steelhead were collected in 22 sites which were electrofished.

Big Creek Tributaries

We attempted to snorkel additional transects in Monumental Creek in August 1983, but turbid water conditions prevented our surveys. Our August 1981 surveys observed densities of 216 steelhead per 100 m² in the lower 3 km of Monumental Creek. We also observed cutthroat, juvenile chinook salmon and mountain whitefish in that section. It is possible that large sediment influx to Mule and Monumental creeks in July 1981 caused some rearing fish to emigrate from the section.

We observed rearing habitat in Monumental Creek from the West Fork Monumental Creek to Big Creek during 1983 spring surveys. The previously described sediment influxes to Mule and Monumental creeks have probably affected the rearing capabilities of Monumental Creek. Embeddedness sampling by Burns (1983) illustrates the extensive sedimentation of the substrate. Invertebrate habitat utilized immediately below Mule Creek was limited to the swiftest portions of the stream (James M. Montgomery, Consulting Engineers 1983). Slower moving areas were completely covered with sediment.

Personnel from Montgomery Engineers conducted electrofishing surveys of eight 60 m transects in Monumental Creek on October 5-6, 1983, between Buck and Mule creeks (James M. Montgomery, Consulting Engineers 1983). A total of 29 juvenile steelhead were captured, which ranged from young-of-the-year (35 to 70 mm) to 215 mm (fork length). Steelhead were present throughout the transects below Century Creek and were most abundant below Mule Creek. Bull trout, juvenile chinook salmon, mountain whitefish and cutthroat were also captured. Surveys in July and August would provide more reliable estimates of maximum rearing densities, since emigration may have begun in October.

Brush Creek

Brush Creek contains scattered rearing areas throughout the area surveyed (mouth to 0.8 km above Horn Creek). A hydropower diversion for the Flying "B" Ranch creates a barrier to migration approximately 1.6 km above the mouth. We captured juvenile steelhead, cutthroat and bull trout below the diversion (Table 9). A small, unscreened irrigation ditch approximately 0.4 km below the diversion may impact migrating fish, in which case, some form of screening would be beneficial (May 1983).

A natural rock barrier exists approximately 0.8 km above the diversion, and we captured rainbow, cutthroat and rainbow x cutthroat trout hybrids between the two barriers (Table 9). Above the barrier we also captured rainbow, cutthroat and hybrids. Cutthroat comprised 20% of the catch above the barrier and 10% below.

Camas Creek

To monitor the annual trend in juvenile steelhead densities within tributaries, we recounted five transects in Camas Creek from 1981 to 1983 (Table 8). Densities of juvenile steelhead remained very similar from 1981 to 1983, ranging from 7.3 to 9.5 fish/100 m². Using a Kruskal-Wallis test, densities of Age 1+ steelhead and total steelhead (all age groups) were not significantly different from 1981 to 1983 (P<0.05).

Juvenile chinook salmon were uncommon in 1981 and abundance increased nearly ten and six times in 1982 and 1983, respectively. Juvenile steelhead comprised 99% of the fish caught in Camas Creek in 1983 (Table 9).

Yellowjacket Creek, the largest tributary to Camas Creek, supports an abundant population of steelhead between the Yellowjacket Mine and Camas Creek. We observed 157 in 5 transects for a density of 9.9 fish per 100 m² (Table 8). Steelhead comprised 99% of the fish captured, followed by cutthroat, bull trout and rainbow x cutthroat trout hybrids (Table 9). Steelhead ranged from 90 to 260 mm (Fig. 13).

Table 9. Numbers of fish sampled by hook-and-line in Middle Fork Salmon River tributaries, July-August, 1983.

Date	Section	Steelhead	Cuttthroat	Bull trout	Hybrid (rainbow x cutthroat)		Mountain whitefish	Chinook salmon	Total game fish
Brush Creek									
28 July	Above diversion	31	8	0	2	0	0	0	41
28 July	Below diversion	18	2	0	1	0	0	0	21
	Totals	49	10	0	3	0	0	0	62
Camas Creek									
20 July	Dry Gulch to Hammer Cr.	63	1	0	0	0	0	0	64
21 July	West Fork	5	0	0	0	0	0	1	6
1 Sept.	Dry Gulch to Hammer Cr.	51	0	0	0	0	0	0	51
	Totals	119	1	0	0	0	0	1	121
Yellow Jacket Creek									
14 July	Mine to trailhead	108	0	1	1	0	0	0	110
19 July	Mouth to trailhead	101	1	0	0	0	0	0	102
	Totals	209	1	1	1	0	0	0	212
Loon Creek									
7 July	Diamond "D" upstream	3	1	2	0	0	0	0	6
6-7 July	East Fork Mayfield Cr.	34	0	8	0	2	0	0	44
	Totals	37	1	10	0	2	0	0	50
Marble Creek									
22 Aug.	Sunnyside-Safety Cr.	0	20	0	0	0	0	0	20

Table 9. Continued

Date	Section	Hybrid					Total
		Steelhead	Cutthroat	Bull trout	(rainbow x cutthroat)	Mountain whitefish	
Marble Creek (cont'd)							
23 Aug.	Safety-Cornish Creek	0	35	0	0	0	35
23 Aug.	Cornish-1 km downstream	7	13	1	0	0	21
23 Aug.	1 km below Cornish-Cottonwood Creek	4	23	1	0	0	28
5-6 Sept.	Mouth-Mitchell Ranch	66	15	0	0	3	86
6 Sept.	Canyon-Birch Creek	9	2	0	0	0	11
6 Sept.	Above Birch Creek	<u>44</u>	<u>16</u>	<u>0</u>	<u>1</u>	<u>3</u>	<u>65</u>
	Totals	130	124	2	1	6	266
Sheep Creek							
29 July	Lower 3.2 km	16	1	0	0	0	17
Sulphur Creek							
29 Aug.	Mouth to North Fork	15	0	0	0	0	18
30 Aug.	In North Fork	0	19	2	0	0	21
31 Aug.	In South Fork	<u>0</u>	<u>23</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>23</u>
	Totals	15	42	2	0	0	62
Wilson Creek							
25-26 July	Above barrier	119	0	0	0	0	119
27-28 July	Below barrier	<u>51</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>51</u>
	Totals	170	0	0	0	0	170
GRAND TOTALS		745	180	15	5	8	960




Figure 13. Length frequency of juvenile steelhead caught in Yellowjacket Creek, July 1983.

We captured juvenile steelhead and chinook salmon in the West Fork Camas Creek up to Pole Creek.

Loon Creek

Transects were surveyed in Loon and Warm Springs creeks in 1981. We surveyed Loon Creek above Mayfield Creek and the East Fork Mayfield Creek in 1983. Both sections contain suitable rearing habitats and we captured steelhead, cutthroat, bull trout and mountain whitefish (Table 9).

Marble Creek

We observed juvenile steelhead in Marble Creek up to Cornish Creek. The section between Big Cottonwood Creek and Sunnyside Creek supported an abundant population of cutthroat (Table 8). Young-of-the-year salmonids were present in all transects, indicating the presence of nearby spawning areas. We also observed cutthroat in both Sunnyside and Cornish creeks (Table 6).

Cutthroat comprised 88% of the fish captured above Cottonwood Creek, followed by steelhead and bull trout (Table 9). Most cutthroat ranged from 60 to 230 mm (Fig. 14).

We were not able to snorkel sections of Marble Creek below Cottonwood Creek due to turbid water conditions. Hook-and-line surveys in August 1982 and September 1983 documented a large abundance of juvenile steelhead below Trail Creek. In 1982, we captured 151 game fish in approximately 20 man hours of angling on Marble Creek below the Mitchell Ranch. Juvenile steelhead comprised 89% of the catch, cutthroat 8% and juvenile chinook salmon 3%. In 1983 we captured 162 game fish on an angling survey between the mouth and Trail Creek. Steelhead comprised 73% of the catch and cutthroat 20%, with juvenile chinook salmon, rainbow x cutthroat trout hybrids and mountain whitefish also present (Table 9).

Juvenile steelhead captured below Trail Creek ranged from 80 to 300 mm (Fig. 14). Cutthroat in the same area ranged to 380 mm, and large cutthroat (>250 mm) were more abundant in sections below, than above, Cottonwood Creek.

In August 1980, Reingold (1981) surveyed the entire length of Marble Creek with hook-and-line gear and also observed cutthroat to be most abundant in upper sections and steelhead in lower areas. He observed juvenile chinook salmon in most pools between Dynamite and Canyon creeks. Six adult chinook salmon and four redds were observed in the same area. Numerous fry and Age 1 fish were observed in the lower 0.4 km of Trail Creek. Idaho Department of Fish and Game biologists have photographs of chinook salmon spawning near the mouth of Sunnyside Creek in 1978.




Figure 14. Length frequency of juvenile steelhead and cutthroat trout caught in Marble Creek, 1982-1983.

Personnel from Montgomery Engineers surveyed sections of Marble Creek on September 16, 1981, September 23, 1982 and October 5, 1983 near Sunnyside Creek and 0.8 km below Safety Creek (James M. Montgomery Consulting Engineers 1983). For the three samplings combined, the consultants captured 66 cutthroat and 1 steelhead near Sunnyside Creek and 70 cutthroat below Safety Creek. Sculpin were abundant at both sites. Fork lengths of cutthroat and steelhead ranged from 30 to 170 mm and 60 to 120 mm, respectively. Benthic macroinvertebrate taxa and densities were also sampled (James M. Montgomery Consulting Engineers 1983). Eleven orders represented by approximately 25 families and at least 40 genera were sampled.

Papoose Creek

A probable migration barrier exists in Papoose Creek approximately 200 m above the mouth. We did not survey fish populations in the stream.

Sheep Creek

Juvenile steelhead were abundant in the lower 3 km of Sheep Creek and we observed 90 in five transects (Table 8). Cutthroat and bull trout were also present.

Sulphur Creek

We observed juvenile steelhead, cutthroat, juvenile chinook salmon and mountain whitefish in Sulphur Creek downstream from North Fork Sulphur Creek (Table 6). Juvenile steelhead were most abundant in the pocket water below the Morgan Ranch. Above that point, we observed or captured only cutthroat and bull trout (Table 9).

An isolated population of cutthroat occurs above a migration barrier, approximately 3 km above the North Fork.

Wilson Creek

A migration barrier occurs in Wilson Creek approximately 0.4 km below the confluence of Alpine Creek. Above the barrier, we observed an abundant population of residualized steelhead or resident rainbow trout and bull trout (Table 6). Below the barrier, we observed steelhead and mountain whitefish. Fish below the barrier had a length frequency distribution similar to those above the barrier (Fig. 15).




Figure 15. Length frequency of juvenile steelhead-rainbow caught in Wilson Creek, above and below a barrier, July 1983.

Life History and Movements

The life history and movements of Middle Fork steelhead are complex and variable. Differences in time of entry into the upper Salmon River, migration speed and seasonal staging are likely influenced by both environmental and genetic factors.

Middle Fork steelhead are summer-run fish (migrating into the Columbia River in summer), which appear to be predominantly "Group B" steelhead. By definition, "Group B" steelhead pass Bonneville Dam after August 25 and a majority are large (exceeding 71 cm) after spending two years in the ocean. Middle Fork steelhead are predominantly large fish, averaging 81 to 86 cm. We captured 170 adult steelhead in the Middle Fork from 1981 to 1983 and 85% exceeded 76 cm (Thurow 1983). Fish ranged up to 99 cm. Female steelhead appeared to be predominant in the early portions of the run, as sex ratios averaged 5.2 females per male in October-November 1981 and 3.3 females per male in March-April 1982.

A portion of the steelhead destined for the Middle Fork ascend the Salmon River in fall, while the remainder over-winter in the Snake River (Mallet 1970). A portion of the run stages in pools below the Middle Fork, while some fish "wander" widely above and below the Middle Fork. Most wild steelhead begin moving above the South Fork Salmon River after mid-September. A segment of the run enters the lower 10 to 15 km of the Middle Fork in fall (Table 10). We observed increasing numbers of steelhead staging in the lower Middle Fork after October 1 (Fig. 16). On August 30, 1982, we observed eight adult steelhead in the lower 0.8 km of the Middle Fork. The fish appeared to be large (>71 cm) wild fish, but due to timing, were obviously not group "B" steelhead. Local anglers recalled that during the 1950's, anglers occasionally caught "large numbers" of steelhead during August in the lower km of the Middle Fork. It is possible that both Group "A" and "B" steelhead spawn within the Middle Fork drainage.

Historically, anglers caught adult steelhead more than 48 km upstream on the Middle Fork in October and November. Between 1981 and 1983, we could not locate any fish more than 14 km up the drainage in fall. Presently only small numbers of steelhead stage in the lower portions of the drainage in fall.

Returns of tagged steelhead illustrate that many of the adult steelhead which currently ascend the Middle Fork in fall do not overwinter there. Instead, they re-enter the main Salmon River, likely with the onset of winter.

In the spring, steelhead begin staging in the lower 0.8 km of the Middle Fork. These fish are extremely vulnerable to angling, and in 1982, three Department personnel captured 33 fish in a single day. Numbers of staging steelhead increased in April, and on April 1, 1983, I observed 72 adult steelhead in the lower 400 m of the Middle Fork (Fig. 16). The spring migration of steelhead into the Middle Fork resulted in large catches near the mouth that were documented during March and April, 1958 to 1962 (Idaho Department of Fish and Game files, Boise).

Middle Fork
adult steelhead
1983

Table 10. Results of spring and fall surveys for adult steelhead in the Middle Fork Salmon River drainage, 1983.

=====

Date	Location	River km	Number steelhead caught or observed
15-17 March	Flying "B" Ranch to Camas Creek	48-57	3 adults, river km 51
4-7 October	Flying "B" Ranch to main Salmon River	0-48	4 adults, river km 1
2-4 November	Big Creek vicinity	26-31	No adults
15-18 November	Flying "B" Ranch to main Salmon River	0-48	2 adults, river km 1




Figure 16. Numbers of adult steelhead observed in the lower 800 m of the Middle Fork Salmon River, 1981-1983.