

UNIVERSITY OF IDAHO
COLLEGE OF AGRICULTURE


EXTENSION DIVISION

E. J. IDDINGS
DIRECTOR

Swine Husbandry In Idaho

By

J. E. NORDBY and E. M. GILDOW



Grand Champion Poland China Boar.
A Sire of Champions.

COOPERATIVE EXTENSION SERVICE IN AGRICULTURE
AND HOME ECONOMICS OF THE STATE OF IDAHO
UNIVERSITY OF IDAHO EXTENSION
DIVISION AND U.S. DEPARTMENT
OF AGRICULTURE
COOPERATING

ANIMAL HUSBANDRY SECTION

Printed and distributed in furtherance of the purposes of the Cooperative
Agricultural Extension Service provided for in Act of
Congress, May 8, 1914.

Table of Contents

Breeding, Feeding and Management

	Page
Type	5
Selecting a Breed	6
Grading up the Herd	6
Selecting the Breeding Herd	6
Food Nutrients for Swine	9
Quality of Nutrients in Feed	11
Feeding Stuffs	12
Forage Crops	14
Hogging-Off Crops	15
Supplements	17
Succulent Feeds	20
Method of Feeding	20
Salt	20
Minerals	21
Hairless Pigs	21
Medical Condiments	21
Water for Swine	22
Housing and Equipment	22
Feeding and Management	33

Swine Diseases, Prevention and Control

Prevalence of Swine Diseases in Idaho	45
Swine Sanitation	45
Advantages of Hog Lot Sanitation	46
Preventing the Introduction of Diseases	47
Elimination of Diseases in the Herd	48
Control of Hog Diseases	49

Swine Husbandry in Idaho

J. E. NORDBY and E. M. GILDOW*

SWINE production is carried on for the most part in conjunction with other farm enterprises. Uniform success in the industry is based upon the efficient utilization of by-products, and upon foraging and "hogging off" plans that are sufficiently flexible to make possible the production of pork with the smallest amount of grain and labor. The swine population is confined largely to the irrigated sections of southern Idaho, and wheat producing areas of northern Idaho (*Fig. 1 and Table I*). Corn, wheat and barley, dairy by-products, soybeans, cull peas, cull potatoes, and an abundance of alfalfa are utilized in the irrigated sections. Successful ration combinations are made from these for all phases of swine production. Wheat, barley, oats, peas, cull potatoes, soybeans and corn in restricted areas, some dairy by-products, alfalfa and other forage crops are used in the northern part of the state.

The attitude of the producer toward the details involved in swine production is one of the most important elements which contributes to success or failure in the enterprise. His livestock-mindedness is an asset that contributes to stability if hog prices and production costs are not in tune. The adaptability of hogs to each individual farm is

TABLE I
Swine on Farms in Idaho by Counties April 1, 1930,
U. S. Agricultural Census, 1930

County	No. Hogs	County	No. Hogs
Ada	12,603	Gem	4,488
Adams	2,855	Gooding	7,618
Bannock	10,441	Idaho	21,010
Bear Lake	1,621	Jefferson	9,901
Benewah	1,946	Jerome	8,044
Bingham	12,983	Kootenai	2,328
Blaine	2,962	Latah	12,134
Boise	913	Lemhi	4,533
Bonner	1,450	Lewis	5,182
Bonneville	17,246	Lincoln	4,246
Boundary	888	Madison	5,331
Butte	3,048	Minidoka	5,182
Camas	1,087	Nez Perce	11,280
Canyon	30,630	Oneida	2,389
Caribou	842	Owyhee	4,572
Cassia	11,258	Payette	5,909
Clark	454	Power	1,361
Clearwater	1,788	Shoshone	236
Custer	1,705	Teton	1,477
Elmore	798	Twin Falls	12,910
Franklin	4,785	Valley	613
Fremont	5,674	Washington	8,893
		Total	267,614

determined by a number of factors. The size of the farm, the availability of suitable forage crops, the possibility of growing crops to be "hogged off," and the amount of crop residues that may be avail-

*Assistant Animal Husbandman, and Veterinarian, respectively, Idaho Agricultural Experiment Station.

Drawings of equipment and bills of materials in this bulletin are by courtesy of Department of Agricultural Engineering, Idaho Agricultural Experiment Station.

able from year to year are important in determining how many hogs to keep. The number may vary some from year to year. In wheat producing areas, where corn is not produced, it is well to keep in mind that the price of hogs does not always vary with the price of wheat. The correlation between the price of wheat and hogs is much less in evidence than it is between corn and hogs. One should work into the business of swine production gradually as this method makes possible a better understanding of the details which solve in a large measure the question of numbers for each farm.

Swine production contributes to the stability of the agricultural prosperity of Idaho in the proportion in which farmers realize the necessity of producing some hogs on each farm adapted to swine production, and not by concentrating the production on a few farms. The former method identifies the swine production enterprise as an adjunct to the general farm production policy, while the latter practice may make it necessary to farm for the sake of the swine enterprise. This often proves disturbing when crop prices are not necessarily correlated with prices of market hogs.

Type

Type in swine is recognized as a combination of those characteristics which contribute to efficiency in the cost of production, and to adaptability in pork trade requirements. Type in swine is determined almost entirely by the producer of market hogs, the breeder of purebred swine, and by the consumer of pork products. The consumer, no doubt, has had a major part in this program and exerts his influence directly through his exacting demands at the meat counter. These demands are reflected to the producer of market hogs through the retailer and packer who pay premiums for hogs that are acceptable in weight, quality and shape, and discount heavily coarse, over-weight hogs. The growing demand for leaner, more tender, juicy, and smaller cuts with less waste has made it necessary for the breeder of purebred swine to supply a type which makes it possible for the producer of market hogs to breed and develop individuals in which there is the closest possible coordination in the characteristics which influence cost of production, efficiency, and consumers' demands.

The breeder of purebred swine is vitally interested in the inherent ability of his hogs to grow rapidly, to be straight and sound in the feet and legs, strong in the back and to have the required depth, thickness and smoothness of body, and quality and style that make them wear and sell well. These rigid requirements will make it possible for the commercial producer to put on the market at six months of age 200-pound firmly finished barrows that have long, deep and smooth sides; thick, full and smooth hams; a moderately wide and gracefully carried back; deep and smooth shoulders; and sound feet and legs. When all of these characteristics are combined in a well-balanced individual with an abundance of quality, the exacting demand of the consumer is likely to be satisfied (*See Fig. 2*).

Selecting a Breed

Personal preference on the part of the breeder is one of the important elements in selecting a breed. He should try to satisfy this desire as far as possible. One usually gets the best results with the breed one likes. Results in the feedlot do not bring out much difference between breeds when the individuals are selected with equal care and given the same attention. Prolific strains that are hardy and have good rustling and feeding qualities should be selected in any breed. The ability of the sow to mother her young is fundamentally important.

Grading Up the Herd

Uniformity in feeder hogs can be encouraged by constantly selecting purebred sires from one breed. When this plan is followed for a



Fig. 2.—Grand Champion pen of barrows, Pacific International. Well finished, high quality, 200-pound barrows.

few years, the grade pigs will be virtually as uniform as good purebreds in type and color. Boars selected for this purpose should come from herds where there is evidence of constructive effort to produce a good uniform type.

Selecting the Breeding Herd

Experimental data have justified the conclusion that the time and cost required to produce a 200-pound market hog decreases as the percentage of pure breeding increases. Obviously, therefore, it is desirable to use very high grade or purebred sows for foundation herds. The price of purebred sows, that are acceptable in type, is in line with commercial values under normal conditions. The breeder of purebred swine is constantly making efforts to maintain a uniform and efficient type in his herd. Sows selected from such a herd will usually prove to be more satisfactory for the production of market

hogs than will grade sows which do not have such a desirable selective heritage to their credit.

The Sow. A brood sow should have considerable length and a uniform balance of length in neck, shoulder, back, and rump. The width of the body should be medium and uniform throughout the entire length. The depth should be emphasized, for upon it depends, in large measure, the usefulness of the sow as a producer and the ability of her pigs to be efficient feeders. Smoothness should be evident in the shoulders, sides and hams, for this is one of the characteristics that contribute to quality. The shoulders should be deep and laid closely at the top. The ham must be deep, full and smooth. The back should have evidence of strength and be gracefully carried




Fig. 3.—Grand Champion Duroc Jersey sow, Pacific International. A deep, smooth, sound individual and a dam of champions.

in a uniform arch. A deep, long, and smooth side is necessary. The joints should be neat, and free from coarseness and the shoulder free from excessive fullness. The width and length of the head must balance. Too much length is undesirable for it is usually associated with a narrow and pointed snout. The ears should be moderate in size, and lightly carried. The eyes should be reasonably prominent. Wrinkles in the forehead of a sow often grow so large that the folds close the eyelids. Neatness and quality about the head are therefore imperative. Sound feet, and legs set squarely under the body are desirable. A brood sow should have all of these characteristics properly coordinated in a nicely balanced and breedy individual (See Fig. 3).

The tendency to be vicious, nervous and irritable may be inherited. A sow having such a tendency should be eliminated from the

herd. The test of a good sow lies in her ability to produce pigs at a profit. The sow should have at least ten normal properly spaced teats. The tendency to produce "blind" teats appears hereditary from data secured at this station. When the end of the teat is telescoped back into the body of the teat it is spoken of as a "blind" teat. (See Fig. 4).




Fig. 4.—A useful sow should be equipped with at least six good teats in each of the two rows. A, good underline; B, sow with three blind teats on each side and in the middle of the row; C, the same sow after nursing a small litter for four weeks. Three teats on each side functioned. (See text for inheritance of this defect.)

The Boar. The boar should always be a creditable purebred individual. His initial cost is very little more than the cost of a grade. The use of purebred boars is so general that it is rather odd to speak of grade boars. A boar with a creditable conformation is more often a good sire than is a boar inferior in individuality. All purebred boars are not good boars.

A boar should, in general, be characterized by ruggedness and masculinity. These should not be emphasized, however, at the expense of quality. Masculinity must not be confused with coarseness and viciousness. A boar may be very high in quality and still be rugged and masculine. Thick, wide and coarse shoulders are often erroneously associated with masculinity in a boar. They are too often the result of hereditary influences and will usually be transmitted to the offspring. It is impossible for a coarse-shouldered pig to develop into a smooth barrow.

Size is difficult to maintain in swine under the conditions that usually prevail in commercial production. It is advisable, therefore, to use a boar that has indications of a large growth heritage. The most economical gains are made when pigs are growing. Some advantage is offered, therefore, by using a boar whose pigs will not slow up their growth too soon.

A boar should have ample length that shows the proper balance in neck, shoulder, back and rump; shortness in the neck and back must be emphasized. The rump should be long and well developed, and the entire top line heavily muscled and supported in a smooth and uniform arch. The loin must be full and smooth. Thin-loined boars often weave when walking. Superior sides are invariably deep with the underline neatly carried. The hams must be smooth, deep, wide, and nicely balanced. The shoulders should be deep, neatly laid at the top and should not carry too heavy a shield. The shield is of service in combat only, hence, is of little use under our domesticated conditions. The bone should be large, but not coarse. The legs must be properly placed, making it possible for the boar to walk straight. The walk of a boar is important and he cannot walk straight if all parts of the body, including the feet and legs, do not properly coordinate. A boar with a scrotal hernia or with only one testicle in the scrotum should not be selected as a sire as both these defects are apparently inherited.

Food Nutrients for Swine

The value of a feed is determined primarily by the amount of digestible nutrients it contains. These nutrients are proteins, carbohydrates, fats, minerals and vitamins. Proteins are necessary for developing muscle. Carbohydrates and fats are essential for providing heat or energy for the body. Minerals contribute to the growth of the bony structure of the body, and to the performance of important body functions. Vitamins are necessary for growth, reproduction, and for the animals' vitality in general.

Inasmuch as the feed cost of producing pork varies from 75 to 85 per cent of the total cost it is obvious that one should give much

attention to feeds. Practically all the feeds used should be home-grown. Purchased feeds should contain as large a percentage as possible of the particular element that is lacking in the home-grown feeds. This element is usually protein. It should also be an animal protein. Some ready mixed feeds may run rather high in protein but the bulk of it may be vegetable proteins that are found in home-grown feeds.

TABLE II
Composition of Some Common Feeds.*

Kind of Feed	Total dry matter in 100 lbs. of feed	Digestible nutrients in 100 pounds of feed				Nutritive ratio of feeds†
		Crude protein	Carbo-hydrates	Fat	Total	
<i>Concentrates</i>						
Barley (Common)	90.7	9.0	66.8	1.6	79.4	1: 7.8
Corn (Dent)	89.5	7.5	67.8	4.6	85.7	1:10.4
Oats	90.8	9.7	52.1	3.8	70.4	1: 6.3
Peas (Field)	90.8	19.0	55.8	0.6	76.2	1: 3.0
Wheat (all analyses)	89.8	9.2	67.5	1.5	80.1	1: 7.7
Beans (Navy)	86.6	18.8	51.3	0.8	71.9	1: 2.8
Fish Meal	89.5	40.1	8.3	58.8	1: 0.5
Linseed Oil Meal (O. P.)	90.9	30.2	32.6	6.7	77.9	1: 1.6
Soybean seed	90.1	33.2	24.7	16.1	94.1	1: 1.8
Tankage (60% protein) ..	92.1	56.2	7.2	71.4	1: 0.3
Tankage (below 40%) ..	90.9	34.3	14.1	66.0	1: 0.9
Wheat bran (all analyses) ..	89.9	12.5	41.6	3.0	60.9	1: 3.9
Wheat Middlings (flour) ..	89.3	15.7	52.8	4.3	78.2	1: 4.0
Wheat Middlings Standard (shorts)	89.5	13.4	46.2	4.3	69.3	1: 4.2
<i>Milk and Milk Products</i>						
Cow's milk	13.6	3.3	4.8	3.6	16.2	1: 3.9
Buttermilk	9.4	3.4	4.9	0.1	8.4	1: 1.5
Buttermilk (dried)	88.3	29.3	41.0	6.2	84.2	1: 1.9
Buttermilk (semi-solid) ..	35.0	12.6	16.7	3.5	36.2	1: 1.9
Skim milk (centrifugal) ..	9.9	3.6	5.1	0.2	9.1	1: 1.5
Skim milk (dried)	95.5	32.5	49.9	1.9	86.7	1: 1.7
Whey	6.6	0.8	4.7	0.3	6.2	1: 6.8
<i>Dried Roughages</i>						
Alfalfa hay (all analyses) ..	91.4	10.6	39.0	0.9	51.6	1: 3.9
Alfalfa leaves	93.4	17.3	35.9	3.0	60.0	1: 2.5
<i>Roots and Tubers</i>						
Beet, sugar	16.4	1.2	12.6	0.1	14.0	1:10.7
Carrot	11.7	1.0	9.1	0.2	10.6	1: 9.6
Potato	21.2	1.1	15.8	0.1	17.1	1:14.5
<i>Green Legumes</i>						
Alfalfa (before bloom)	19.9	3.5	7.5	0.3	11.7	1: 2.3
Peas (Canadian field)	16.6	2.9	7.1	0.3	10.7	1: 2.7
Soybean (in bloom)	20.8	3.0	8.5	0.3	12.2	1: 3.1
<i>Green Fodder from the Smaller Cereals</i>						
Barley fodder	23.2	2.3	11.5	0.4	14.7	1: 5.4
Oat fodder (8 in. high) ..	13.0	3.4	4.1	0.5	8.6	1: 1.5
Rye fodder (5 in. high) ..	18.1	5.1	6.2	0.7	12.9	1: 1.5
Wheat fodder (5 in. high) ..	24.2	5.1	10.3	0.5	16.5	1: 2.2
<i>Miscellaneous</i>						
Apple	18.2	0.4	15.6	0.2	16.4	1:40.0
Apple pomace	23.3	1.2	15.6	0.8	18.6	1:14.5
Pumpkin (field)	8.3	1.1	4.5	0.5	6.7	1: 5.1
Rape	16.7	2.6	10.0	0.3	13.3	1: 4.1

*From Henry and Morrison's "Feeds and Feeding."

†This ratio is obtained by adding to the carbohydrates 2.25 times the amount of fat and dividing the sum by the amount of crude digestible protein.

The following recommendations for nutritive ratios of the grain feed are made merely as a general guide and are approximately correct when 50 per cent of the protein supplements used are from an animal source:

	Nutritive Ratio
Growing pigs on a legume pasture	1:6.5
Fattening pigs in a dry lot	1:5.5
Brood sows in winter, getting alfalfa hay	1:6.0
Brood sows during suckling period on legume pasture ..	1:6.5

Home-grown feeds vary considerably in their nutritive value. These variations are shown in Table II. In this table are found differences in the protein, carbohydrate and fat content of a number of feeds that are used in swine production. In the last column to the right is found the nutritive ratio of the feeds. Since feeds are usually compared on the basis of their nutritive ratio it is well to know just what it represents. In the table we find the nutritive ratio of barley to be 1:7.8 which means that if we give the value of 1 to the amount of digestible crude protein (9.0) in 100 pounds of barley the value of carbohydrates (66.8) plus the value of the fat (1.6) would be 7.8. The carbohydrates and fats are combined because they serve essentially the same purpose. One pound of fat, however, has a value equal to 2.25 pounds of carbohydrates. In order to combine these two, therefore, they must be reduced to a common basis. This is done by multiplying 1.6 (per cent of fat in barley) by 2.25 which gives 3.6. When we add 3.6 (the carbohydrate equivalent of 1.6 pounds of fat) to 66.8, the carbohydrates in barley, we get 70.4. When we divide 70.4 by 9.0 (crude protein) we find that 70.4 is 7.8 times as large as 9.0. The nutritive ratio is, therefore, expressed as 1 to 7.8.

Nutritive Ratio of Mixtures. It is not quite so simple to find the nutritive ratio of a combination of feeds, at least when they contribute different amounts of the mixture. When any feed in a mixture makes up less than 100 pounds, the figures for the digestible protein, carbohydrates and fat in the table, which are given for 100 pounds of feed, must be divided by 100 and then multiplied by the pounds of this feed present in the mixture. Find for example, the nutritive ratio of the following mixture:

	Crude Protein	Carbo- hydrates	Fat
Barley, 100 lbs.	9.00	66.80	1.60
Wheat, 100 lbs.	9.20	67.50	1.50
Tankage, 16 lbs.	8.99		1.15
Alfalfa leaves, 10 lbs.	1.73	3.59	.30
	<hr/>	<hr/>	<hr/>
	28.92	137.89	4.55

Applying the formula: Carbohydrates plus 2.25 times the fat, divided by the crude protein or $137.89 \text{ plus } (4.55 \times 2.25) \div 28.92$ or 5.1. Therefore, the nutritive ration of this feed ration is 1:5.1.

Quality of Nutrients in Feed

There is considerable variation in the quality of nutrients in feed that is not recognized in Table II. This is stressed more in connection with the proteins, minerals and vitamins. The amount of protein may be the same in two feeds, yet the proteins in one may be far less complete than the proteins in the other. Proteins are made up of many different parts, some of which are essential to growth and others necessary for maintenance. Those elements in the proteins necessary for growth for example, are largely absent in some feeds even though these feeds may be high in total proteins. One

should, therefore, make up a ration of different kinds of feeds so as to more nearly insure a balanced supply of proteins. Proteins from an animal source, such as tankage and skim milk are important for this reason.

Feeding Stuffs

The large variety of feeding stuffs produced in Idaho makes it comparatively convenient to make up rations that are admirably adapted for the production of pork.

Corn. Corn matures satisfactorily in a number of the irrigated sections in the state, and is also produced in limited sections in the Clearwater drainage area of Northern Idaho. Much of the corn produced in the state is fed to swine. Corn that is properly matured here compares favorably with eastern corn as a feed for swine. A protein supplement should be fed with corn. When the pigs are on a legume pasture and corn, it is helpful to feed one pound of skim milk for each pound of corn, or give them five pounds of tankage with each 100 pounds of corn.

Barley. Barley is used extensively in pork production in Idaho. Its feeding value is about 8 per cent less than the feeding value of No. 2 corn (Table III). It should be ground or rolled for best results. Eighty-five-pound fall pigs produced 100 pounds of gain with 306 pounds of barley, 131 pounds of peas and 25 pounds of alfalfa hay, gaining at the rate of 1.2 pounds per day. In other tests at this station, 100 pounds of pork have been produced in the dry lot by pigs weighing 84 pounds at the beginning of the test with 430 pounds of barley, 20 pounds of 60 per cent digester tankage, and 20 pounds of alfalfa leaves. In the same trials it required 580 pounds of barley when fed alone to produce 100 pounds of gain.

TABLE III

Comparative Value of Fattening Feeds for Hogs When All Feeds Noted Are Properly Supplemented.

(Ground barley used as a standard for comparison.)

Feed	Approximate comparative value of 100 pounds when ground barley is worth									
	Per cent*	Dollars								
Ground barley	100	\$0.60	\$0.70	\$0.80	\$0.90	\$1.00	\$1.10	\$1.20	\$1.30	\$1.40
Ground corn	109	.65	.76	.87	.98	1.09	1.20	1.31	1.42	1.53
Ground wheat	114	.68	.80	.91	1.03	1.14	1.25	1.37	1.48	1.60
Ground oats	82†	.49	.57	.66	.74	.82	.90	.98	1.07	1.15
Wheat middlings	90†	.54	.63	.72	.81	.90	.99	1.08	1.17	1.26

*Barley is used as a standard for comparison. When ground barley is given a value of 100 per cent, corn has a value of 109 per cent and so on for the other feeds. The price comparisons are based upon the values in the per cent column when barley has the price value indicated. There is obviously some variation in actual feeding operations from the comparative per cent values assigned. These values, however, will serve as an approximate guide.

†Value of ground oats when fed not to exceed one-third of the grain ration. Wheat middlings are, as a rule, not fed as a fattening ration.

and when 60 per cent digester tankage was fed with barley, it required 446 pounds of barley and 34 pounds of digester tankage. The pigs fed barley alone gained .84 pounds each day, those fed tankage and barley 1.3 pounds per day, and those fed alfalfa leaves, barley,

and tankage 1.45 pounds per day. Another lot gained 100 pounds on 330 pounds of barley and 142 pounds of peas at the rate of 1.08 pounds per day.

Wheat. It is not good practice to feed wheat alone. It should be fed ground. Soaking ground wheat scarcely pays for the work involved. It is not profitable to feed whole wheat either soaked or dry. Grinding wheat increases its value from 15 to 20 per cent over whole wheat, and increases the daily gains in hogs from 10 to 15 per cent. Wheat when ground is worth from 3 to 5 per cent more than No. 2 corn. Spring pigs weighing 52 pounds gained, on a full grain feed, 1.45 pounds per day on alfalfa forage, and required 348 pounds of wheat and 27 pounds of tankage for each 100 pounds gain. Another lot gained on a limited ration 1.28 pounds per day on alfalfa forage requiring 338 pounds of wheat and 22 pounds of tankage. Approximately 550 pounds of wheat are required to produce 100 pounds of gain in dry lot feeding of fall pigs when wheat alone is fed. When one pound of tankage, fish meal or its equivalent in skimmilk, etc., is fed with 12 to 14 parts of wheat, the wheat requirement for 100 pounds of gain is about 425 pounds. The daily gains also will be larger when a protein supplement is fed.

Satisfactory gains have been secured at this station when wheat has been supplemented with cracked Canadian field peas. One-hundred-fifty-pound fall pigs gained 1.83 pounds per day for 42 days, requiring 396 pounds of wheat and 46 pounds of cracked peas for each 100 pounds gain. Wheat may be fed to all classes of swine when properly balanced with protein supplements. It is not desirable to supplement wheat with only wheat by-products.

Peas. When peas are harvested they are usually too expensive to feed to hogs. Pigs frequently go off feed and make unsatisfactory gains when fed peas alone in the dry lot. In tests at this station with 68-pound pigs fed cracked peas, gains of .76 pound per day were made for 175 days requiring 445 pounds of peas for each 100 pounds of gain. Where peas constitute the only ration for brood sows during the period of pregnancy the pigs farrowed by these sows averaged, in tests at this station, one-half pound less and were less thrifty than pigs farrowed by sows fed barley, cracked peas, and alfalfa hay.

Pea Screenings. Pea screenings can be utilized to advantage. They vary so much in composition, however, that it is difficult to give pea screenings even an approximate value.

Oats. The oats produced in this climate are usually very plump and it is very doubtful if any grain is superior to oats for growing pigs. Oats are likewise very useful for brood sows during pregnancy and as part of the ration during the suckling period.

Oats should not comprise more than one-fourth to one-third of the ration for finishing hogs, they must be ground for best results. If oats are selling for a higher price than other grains, less oats may be fed. It has not proven generally satisfactory in Idaho to feed oats in large quantities in feed lot rations (*Table III*).

Soybeans. A few areas in the state are well adapted to the production of soybeans. Corn and soybeans require about the same climatic conditions for the best yields. Soybeans are high in protein and low-melting oil and should not constitute the entire ration. A large proportion of soybeans in a ration will result in soft pork. It is likely that soybeans will be used primarily as a forage crop in Idaho as they are ready to be pastured when other green feed is scarce. When mature soybeans are hogged-off, the hogs should be fed grain or allowed the run of a cornfield. They are not so palatable for small pigs as for pigs that weigh 100 pounds or more. Soybeans may be fed as a protein supplement up to 10 or 12 per cent of the ration. Other supplements should also be used. Soybeans may also be used for brood sows or for developing breeding hogs.

Beans. When cull table beans are cooked in water to which a small amount of salt has been added, they are satisfactory for swine. When cooked, beans are fed in equal amounts with one of the standard concentrate feeds such as barley, corn or wheat it will require about 425 pounds of the mixture to produce 100 pounds of gain. When beans are fed alone the daily gains are not so satisfactory as when beans are combined with a grain feed. Beans should be cooked before they are fed to swine.

Forage Crops

The successful production of swine in Idaho depends in large part upon the utilization of forage crops. Forage crops not only supply comparatively cheap feed, but they keep the hogs in better physical condition and are indispensable in following sanitation requirements.

When hogs below 200 pounds in weight are on a full grain ration and alfalfa forage, the forage will replace from 15 to 20 per cent of the grain requirements. The forage will replace more grain if the hogs are getting less than a full feed of grain. Mature brood sows that are not suckling pigs will, as a rule, stay in good shape on alfalfa forage when fed even less than one pound of grain per day for each 100 pounds of live weight.

It is advisable, in choosing forage crops, to select those that are adapted to local soil and climatic conditions. Forage crops should also be selected on the basis of cost of production, palatability, yield, succulence, length of grazing period, and ability to endure tramping. They should, when possible, be leguminous in character.

Alfalfa. Alfalfa is the most desirable forage crop for swine. In common with other forage crops, it grows better when it is not pastured too closely. It is a good policy to provide enough acreage so that a crop of hay may be removed during the forage season.

Bluegrass, and other Perennial Grasses. Bluegrass, when properly managed, is used with some success in the irrigated sections of the state. It cannot compete successfully with the legumes in the non-irrigated sections.

Other perennial grasses are not so desirable, as they lack the palatability that is so characteristic of the legumes.

Rye. The value of rye lies in its ability to furnish succulence when the other forage crops are scarce. It does not supply a large amount of forage, but it is utilized to advantage late in the fall and early in the spring.

Wheat. Winter wheat seeded in the spring may be used as pasture the first summer, primarily in areas outside the alfalfa producing sections. It is often foraged the second year until it begins to joint. It should never be grazed closely. Twelve pigs weighing 46 pounds each were turned into a one-acre field of winter wheat May 20, at this Station. The wheat had been seeded in October of the previous year. The pigs were fed two pounds of grain per day for each 100 pounds of live weight, or about one-half as much as they would eat, and gained .81 pound a day for 62 days. They required 257 pounds of wheat and 13 pounds of tankage (60 per cent) for each 100 pounds of gain. A similar lot for the same period on one acre of wheat gained 1.03 pounds per day on a full grain ration of wheat and tankage and required 358 pounds of wheat and 18.5 pounds of tankage for each 100 pounds of gain. Some of the wheat in these lots matured and was hogged-off by the pigs during the test.

Oats. Oats serve only as an emergency forage crop. They are used, however, more satisfactorily in combination with peas. A mixture of 40 pounds of oats and 50 pounds of peas per acre under favorable conditions will produce a fairly satisfactory short period forage crop.

Peas. Peas alone are used most successfully as a hogging-off crop. The yield of forage is comparatively small as peas do not do so well when pigs are constantly picking at them, often pulling up entire plants. As a forage they are successfully used in combination with oats. (See oats above). Peas have no significant advantage over alfalfa.

Soybeans. Hogs are usually turned into the soybeans about a week or ten days after they are in full bloom if one desires to take advantage of the leaves. Soybeans should be grazed lightly at this stage of development if one is interested in having a large per cent of the pods fill.

Hogging-Off Crops

Corn. Quite a large percentage of the corn that is fed to swine in Idaho is hogged-off. Four thousand three hundred acres were harvested by hogs in Idaho in 1929. When hogging-off corn the hogs should be allowed access to alfalfa pasture or some other green feed. The daily gains may be increased if a small allowance of wheat middlings or tankage is supplied.

Peas. An average of 406 pounds of pork per acre of peas hogged-off has been produced in tests over a period of years at this Station. The yields of pork per acre ranged from 250 to 650 pounds. Pigs that are allowed one-half ration of ground barley when they are hogging-off peas gain more rapidly and the carrying capacity of the pea acreage is increased.

Under irrigated conditions in this state a demonstration with 77 pigs turned into an eleven acre field of peas resulted in daily gains of 1.4 pounds per day, for 94 days, and a total production of 824.8 pounds of pork per acre. The pigs that are being finished for market should not be required to clean the fields. This can be done by the stocker pigs and breeding herd.

A pea variety mixture for hogging-off is good in theory only if there is a difference in the time of maturity of the varieties in the mixture. If the hogs are turned in when the earliest variety is ready, the later varieties will be tramped, the effect of which reduces significantly their yield. It is a better practice to use the varieties in separate lots and turn in when each has reached the proper stage of maturity. Peas are ready to be hogged-off when the first pods begin to ripen. Early varieties afford a greater possibility of finishing the hogs in the field for the early market.



Fig. 5.—A good job of hogging off wheat.

Unthreshed pea vines are often harvested for winter feeding. This eliminates the expense of threshing. The vines should be stored under shelter. They should be fed on feeding floors during wet weather.

Wheat. Wheat may be hogged-off with success in dry seasons. The hogs are turned in about as early as the wheat is ready to be cut with the binder. This method appears to be successful enough to merit consideration. Pigs which have been full-fed during the summer do not adapt themselves so readily to hogging-off wheat as those that have had limited rations on forage. Obviously, smooth headed varieties of wheat are more suitable for hogging-off. Pigs that are to be finished should not be expected to clean the fields. This may be done by the breeding herd or stockers (*Fig. 5*).

Soybeans. Soybeans may be hogged-off in conjunction with standing corn. Grain should be fed when hogs are on soybeans, as

they are too high in protein and low-melting oil to be fed alone. The grain may be supplied in a self-feeder or be hand fed.

Supplements

As a general rule pigs that have access to a good legume pasture, preferably alfalfa or clover, when on full feed, require only about one-half as much concentrated protein supplement as they require in the dry lot. If they are not on full feed, they will consume more forage and will require only from one-third to one-half the amount of protein supplement consumed by those in the dry lot. The larger and more mature the hog, the more forage it will consume, consequently less protein supplement will be required. Pigs that are developed to market weight while on a good legume forage require from 15 to 20 per cent less grain for each 100 pounds gain than when in the dry lot. Experiment station workers in general favor protein supplements made up from two or more sources, at least one of which should be from an animal source. Such a plan usually increases the palatability of a ration, encourages greater consumption and more rapid gains.

More rapid gains as well as more economical gains result when the supplements are properly used with grain. The grains are deficient in proteins, minerals, and vitamins—elements which can in large part be supplied by the use of leguminous forages. For best results, protein supplements from an animal source, such as the dairy by-products, tankage and fish meal should be used, as they supply the elements in which the grains are deficient.

Skim milk. Skim milk is very valuable in all phases of swine production. The protein in milk is complete and easily digested. In pasture, pigs utilize skim milk to good advantage if they are fed one pound for each pound of grain when on a full grain ration. In the dry lot, two pounds may be fed for each pound of grain. The value of skim milk in the ration decreases when it is fed in larger quantities. If fed in the amounts indicated, it will require from 400 to 450 pounds of skim milk to replace 100 pounds of grain. When skim milk is available on the farm it may, of course, be fed in larger quantities.

Whey. Whey has a little less than one-half the value of skim milk. Therefore, the cost of handling per unit of protein is materially increased over the cost of handling skim milk. The protein content of whey is low, hence it is difficult to balance the grain ration when it is fed as the only supplement.

Buttermilk. Buttermilk has a value comparable with skim milk, and may be fed in the same amounts. Unless the cream from which creamery buttermilk is made has been pasteurized, it should be heated to 180° F. to help avoid the spreading of diseases, especially tuberculosis.

Dried Buttermilk. Dried buttermilk is a very satisfactory feed for all classes of swine. As a rule it contains less than 40 per cent protein and on the basis of experimental tests it should sell for a little less than 60 per cent digester tankage when used as a swine feed.

Semi-solid Buttermilk. This product contains only from 13 to 15 per cent protein and a little over 60 per cent water. On the average, its value is approximately two-thirds that of 60 per cent tankage. To have this value, however, it must be fed sparingly with other cheaper supplements that make up the bulk of the protein supplements such as tankage.

Tankage. The meat packing industry manufactures different grades of by-products in the form of tankage, often referred to as meat meal, that are valuable in pork production. The per cent of protein in the various grades of tankage should not constitute the sole basis of making comparisons as some of the lower grades are higher in fat. Since tankage is purchased primarily for its protein content, it is generally advisable to buy the grades that contain the highest per cent of protein. One part of tankage and 10 parts of corn or 12 to 14 parts of wheat or barley will, as a rule, give excellent results when fed to hogs in the dry lot. The tankage may be reduced to about one-third or one-half this amount when the hogs have access to high quality alfalfa hay. Investigations point favorably to the use of more than one protein supplement in a ration.

Fish Meal. The composition of fish meal is similar to that of digester tankage. Some tests indicate that high-grade fish meal is about equal to tankage, and others that it is slightly superior to tankage. It can be fed in the same proportion as tankage.

Wheat By-products. Wheat by-products commonly available are bran, shorts, and flour middlings. They vary considerably in composition. In some cases bran is run with the shorts and at times ground screenings may also be included, especially with the middlings. It is difficult therefore, to develop a definite guide for intelligently selecting one or the other of these by-products.

The by-products of wheat are used most satisfactorily in conjunction with other supplements richer in protein, as a basis for slops. Wheat middlings, when substituted for the grain protein of a ration, are worth about 20 per cent less than wheat when fed to swine in the feed lot. While these by-products are high in protein, they do not supplement the proteins in grains satisfactorily, unless the pigs are on legume pasture and are also receiving a protein supplement in the form of tankage, fish meal, or dairy by-products. Wheat bran is too bulky and laxative for little pigs, but is useful for brood sows in limited quantities before farrowing. After farrowing, shorts or middlings are more commonly used. Wheat by-products should not be used alone as supplements for wheat.

The value of supplements is generally calculated on the basis of the amount of grain which they will replace in producing 100 pounds of pork and by the increased rate of daily gain in the hogs. One hundred pounds of tankage will replace from 400 to 450 pounds of grain when digester tankage is fed to the extent of balancing wheat or barley in the dry lot for fattening hogs. It may prove advisable to use less tankage unless one is interested in the maximum rate of

gain and if 400 to 450 pounds of wheat will not buy 100 pounds of tankage.

Peas. Peas may be considered a protein supplement when they are too high in price to feed in larger quantities. With one part of peas and six parts of ground wheat, pigs weighing 150 pounds gained 1.83 pounds per day for 42 days, requiring 442 pounds of feed for 100 pounds of gain. These pigs dressed 79.5 per cent. Peas, however, should not be considered a sufficient supplement for more extended periods as work at this Station points to disturbances arising in the form of lame backs and paralytic rear quarters when 68 pound pigs were fed peas only for 120 days. These difficulties prevailed, although not so generally, in a similar lot fed a mixture of one part peas and 2.5 parts barley, but did not occur in lots fed peas, or peas and barley when a mineral mixture of steamed bone meal 30 parts, finely ground limestone 30 parts, and common salt 20 parts was supplied in the grain ration.

Linseed Oil Meal. About twice as much linseed oil meal as digester tankage is required to balance a ration of wheat or barley. A mixture of two parts of tankage, one part of oil-meal and one part of cut alfalfa hay, leaves or meal is a good protein supplement with corn, wheat or barley. Ten pounds of this mixture may be fed with 100 pounds of barley or wheat.

Soybeans. Harvested soybeans may be fed as a supplement. From 10 to 12 per cent of soybeans may be fed with corn, wheat or barley. Some skimmilk or tankage should be added to the ration for best results.

Alfalfa Hay. It is advisable to use alfalfa leaves or hay in swine feeding. Alfalfa should not be fed to exceed 5 per cent of the ration if rapid gains are required. More may be used when large daily gains are considered not so important. It is profitable, with normal hay prices, to let the hogs work over alfalfa hay that has not been chopped. This method saves the expense of chopping the hay and the hogs will not eat so much of the bulky fibrous stems as they are forced to eat when cut hay is mixed with the grain. Alfalfa should be made available for swine in one form or another throughout the year in regions where it can be produced.

Commercial Feeds. A number of commercial feeds are on the market that carry a moderate amount of protein and are designed to be used in rather large quantities with the cereals. The manufacturer usually advises the use of his product up to one-fourth or one-third of the entire ration. The protein content of these feeds generally is, first of all, too low—the farmer having to buy too many pounds in order to get enough protein, and, furthermore, the protein present is largely of plant origin, the kind which the farmer can produce on his own farm. Purchased supplements should be of animal origin as it is possible to produce sufficient plant protein to satisfactorily produce swine wherever legumes can be grown.

Succulent Feeds

Potatoes. About 425 pounds of cooked potatoes are required to replace 100 pounds of grain when not more than three or four pounds of cooked potatoes are fed with each pound of grain. If two pounds of cooked potatoes are fed with each pound of grain, 350 pounds of potatoes will have the value of 100 pounds of grain. The grain should not be cooked, but may be mixed with the potatoes after they are cooked and cooled. Raw potatoes are only from one-half to two-thirds as efficient as cooked potatoes as a feed for swine.

Roots. Roots are palatable, succulent, and laxative and therefore have a place in brood sow rations. They should be chopped before they are fed to swine. From 400 to 450 pounds of chopped sugar beets will replace 100 pounds of grain when two or three pounds of beets are fed with each pound of grain.

Pumpkins. Pumpkins are fed to swine with success. They are fed raw to the best advantage. The seeds should not be removed when feeding pumpkins, nor should large amounts of seeds be fed alone. The proportion of pumpkins to grain fed for good results should be about two or three pounds of pumpkins to one pound of grain.

Silage... The large amount of fiber in silage makes it undesirable as a feed for hogs. The value in silage lies primarily in the amount of corn or grain which it contains.

Artichokes. This crop does not successfully compete with other crops in the state. They seem adapted, however, to some of the cut-over lands.

Methods of Feeding

Hand Feeding. A number of points must be observed when hand feeding is done. It is better to under-feed a little than to over-feed. Enough trough space should be provided to give all hogs an equal chance when limited rations are fed. This is to be especially emphasized if the hogs are of different size. The feeding should be done at regular intervals and all troughs and floors kept clean so as to maintain the appetite of the pigs.

Self-feeder. The self-feeder method may be used when hogs are on a full grain feed. With this method the hogs have their freedom with respect to the time and amount they eat. Self-feeders should be examined each day to make sure the grain is feeding properly. Self-fed hogs eat more and gain faster, as a rule, than hand-fed hogs; but require about the same amount of grain for the gains made. The self-feeder method is a labor-saver.

A good quality, palatable grain mixture should be supplied in the self-feeder if the protein supplements are fed in a separate compartment or the hogs may eat more than necessary of the more expensive supplements which are, as a rule, palatable.

Salt

All classes of hogs should have salt. It is possibly the only mineral that it is necessary to provide throughout the year when legume

forages and legume hay, dairy products, and tankage are available. Free access to salt when pigs are not accustomed to it, or feeding an overdose often proves fatal (*See treatment for scours under "Swine Diseases" in this Bulletin*). Salt may be fed to the extent of one-half of a pound mixed with each 100 pounds of grain. Larger amounts often cause pigs to scour. If a mineral mixture is fed, the salt may be included. Salt may also be fed with wood ashes. The latter is desirable in getting swine accustomed to salt if later they are to be given free access to it.

Minerals

The common grains are deficient in minerals for hogs. It is common for hogs that are fed only wheat and barley to break down with posterior paralysis, commonly called rickets. The deficiency of minerals often brings about a condition of extreme pains which are plainly evident when the hog moves around. In a number of cases hogs fed the grains alone break down on their way to market. Brood sows that do not get enough calcium fail to produce strong pigs and when they are required to produce litters twice a year, often break down with posterior paralysis.

Protein supplements, such as tankage, fish meal, skim milk and alfalfa, that are commonly used to balance the protein requirement, will generally correct the calcium deficiency in the grains when fed in the amounts recommended for balancing the ration (*see supplements*.) Alfalfa should be supplied throughout the year. It is high in protein, and contains necessary vitamins.

It is probable that mineral mixtures are not so essential if one of the protein supplements mentioned above and alfalfa are fed. However, if it is the desire to feed minerals it is likely that a mixture composed of equal parts of common salt, finely ground limestone, or lime (not quicklime) and steamed bone meal will prove adequate for general conditions in this state. This mixture may be self-fed in a box after the pigs are accustomed to it, or one and one-half pounds of the mixture may be fed with each 100 pounds of grain.

Hairless Pigs

There are a few restricted areas in the state in which hairlessness is found in pigs. In such areas one-third ounce of potassium iodide may be added to each 100 pounds of the limestone, bone meal, salt formula mentioned above. This may be self-fed after pigs become accustomed to it. One ounce of potassium iodide may be added to 100 pounds of common salt and one-half pound of this mixture may be fed with each 100 pounds of feed.

Medicinal Condiments

It is better to give specific worm remedies than to feed complex mineral mixtures that are designed to rid pigs of these parasites. Specific drugs are in the end cheaper and more effective (*See round worms under Swine Diseases*).

Water for Swine

Clean water for swine is as important as good feed. Fresh water should be supplied at least three times daily. If clean running water or an automatic watering device is not available, the water should be made available in troughs that are kept clean. Troughs with cross bars over the top are best for this purpose (*Figs. 11 and 12*).




Fig. 6.—Housing combination in use at Idaho Experiment Station, Moscow. Individual houses below. Central house above which accommodates 20 brood sows in individual pens.

Housing and Equipment

Swine are not able to adjust themselves properly to extreme and sudden changes in climate unless they are adequately equipped with satisfactory shelter. Shelter for swine need not be expensive to be efficient. Houses should be warm, well ventilated, convenient and well lighted. They should also be durable, and it costs little more to make them pleasing in appearance.

The field or colony house and the central house are both used successfully in Idaho (*Fig. 6*). The "A" type field house (*Fig. 7*) may be made very cheaply and will accommodate one sow and a

litter. The one-way roof field house (Fig. 8) affords more room than the "A" house. This house will accommodate two or three sows with their litters after they are trained to stay together. Farrowing rails may be used conveniently in this type of house (Fig. 17). It is very satisfactory for winter management. It can be moved easily. Ample ventilation is provided in both types of houses and in very cold weather a burlap blanket suspended in the doorway helps keep out the chill and allows the hogs to walk in and out at




Fig. 7.—"A" type field house. Plan (A) with rustic siding, roof and ends. Plan (B) with board and battens (Specifications for all types of equipment discussed in this bulletin available through Department of Agricultural Engineering of this Station).




Fig. 8.—One-way field house. A satisfactory type when more room is desired than is available in the "A" type.




Fig. 9.—Portable loading chute.




Fig. 10.—Shipping crate. The dimensions specified for the various weights are satisfactory under normal conditions.

will. A board tacked on the bottom of the burlap blanket will hold it in place. The house should be located with the door on the lee side and with the windows toward the sun as much as possible in the winter. The window side of the house may be turned toward the north in the summer. Field houses are very practical for carrying out the management details in a sanitation program.

Central houses should be carefully located. These are more expensive than the colony types, but are more durable and make




Fig. 11.—Conventional "V" type trough with reinforced ends.




Fig. 12.—A satisfactory bevel box-type trough with reinforced ends.

possible more systematic ventilation, and add to the convenience of the feeding. They are not practical for small herds. With central houses, as a rule, more fencing is required for lanes to outlying pasture lots. Fire risks are greater with central houses.

The individual and central type houses should not necessarily be considered as separate systems.

The two types can be combined when a sufficient number of hogs are produced to warrant the expenditure. (*Plans and specifications for central houses are available through Department of Agricultural Engineering, Idaho Agricultural Experiment Station.*)




Fig. 13.—Convenient castrating crate. By means of this crate the pig can be kept clean during the operation.

satisfactory support is shown in this plan. This support is adequate so long as it has good footing. Whenever it is practical the chute may rest on the wagon or truck floor.

Troughs. This type of equipment is probably the simplest and there are many satisfactory types in use. Troughs are expensive equipment if they are not efficiently made of durable materials. It will be observed that two-inch material is recommended for the troughs in Figs. 11 and 12. Notice also that the ends of each trough carry a two-inch reinforcement that is sawed to fit accurately the inside shape of the trough. This materially strengthens the trough by making it possible to drive the nails into the side pieces in two directions. With this end reinforcement, the trough will not leak so readily. The cross braces can obviously be fastened in different ways. One should, however, not cut into the side pieces of the trough.

Shipping Crates. Shipping crates should be made of light but durable materials. The dimensions indicated in Fig. 10 are satisfactory under normal conditions. When building crates it is well to keep in mind that hogs of the same weight differ considerably in length, width, and height.

Castrating Crate. A castrating crate is of course not an absolutely essential piece of equipment. However, many are in use on farms because they make the work of castrating pigs much easier and also make it possible to keep the pig out of the dirt when the operation is performed. The disinfectant can be kept in a pan on the 1" x 12" shelf immediately under the trough. It is common practice

Loading Chute. When a portable chute is desired, the plan in Fig. 9 may be used. This chute is designed to endure considerable wear and will stand up even though there is appreciable crowding when the hogs are passing up the chute. Portable chutes of this kind must obviously be supported at the wagon end. A




Fig. 14.—Lifting crate.

on many farms to castrate the pig crop on a rainy day when field work has to be temporarily suspended. Under such conditions, the opportunity of doing the castrating job off the wet ground is appreciated. The crate in Fig. 13 suggests one plan that is practical.




Fig. 15.—An inexpensive shade which is practical when natural shade is not available.

Lifting Crate. This crate is used primarily when it becomes necessary to trim the feet of hogs. This phase of management is of course not new to breeders of purebred swine who exhibit their hogs. When the claws on heavy hogs grow out too long, they are likely to break off causing severe lameness, and it may become necessary to confine them in a lifting crate to make it possible to trim the feet properly (Fig. 14).

Wallows. Wallows usually work out better in theory than in practice. They are a source of disease contamination if they are not kept clean. It is practically impossible to keep them sanitary. Protection from excessive heat should be provided by the use of natural shade and running water when that is available.

Shade. Natural shade is desirable. If that is not available, a temporary shade (Fig. 15) may be provided. This should be located where the air is likely to move freely.

TABLE IV

"A" Type Portable Hog House—Building Material for Plan "A" (with rustic siding, roof and end).

Material as Bought			Material as Used		
No. of Pieces	Size of Stock (inches)	Length (feet)	No. of Pieces	Length (feet)	Use
2	4x6	10'	2	10'	Skids
5	4x4	8'	5	8'	Joist
1	4x4	6'	2	3'	Runway joist
12	2x4	8'	12	8'	Plates and rafters
4	2x4	8'	4	2'8"	Studs
			4	5'2"	
1	2x4	8'	1	8'	Ridge pole
1	2x4	14'	1	2'6"	Door lintel
			2	2'	Window
			4	1'8"	Ventilator
1	1x4	10'	2	3'2"	Window trim
			2	1'8"	
1	1x4	8'	2	2'2"	Window casing
			2	1'6"	
4	1x4	8'	4	7'8"	Corner trim
1	1x4	10'	2	3'2"	Door casing
			1	2'6"	
1	1x4	12'	2	3'9"	Door trim
			1	3'2"	
1	1x4	8'	4	1'	Ventilator trim
			2	1'8"	
2	1x6	10'	2	9'	Ridge boards
2	1x4	10'	6	2'4"	Door hurdle
			2	2'10"	
7	2x12	8'	7	8'	Floor
1	2x6	8'	1	8'	Floor
1	2x12	8'	3	2'6"	Floor of runway

40 Bd. Ft. 1"x6"x10'
200 Bd. Ft. 1"x6"x 8'

Summary				Miscellaneous
No. of Pieces	Size of Stock (inches)	Length (feet)	Board Feet	
2	4x6	10	40	
5	4x4	8	53	4— $\frac{1}{2}$ "x10" bolts and washers
1	4x4	6	8	2—8" strap hinges
7	2x12	8	128	2—4" T hinges
1	2x6	8	8	3—3" gate hooks and eyes
17	2x4	8	91	2—4" gate hooks and eyes
1	2x4	14	9	3' wire chain
3	1x4	10	10	6'—1"x $\frac{1}{2}$ " strap iron
1	1x4	12	4	4 lb. 20d nails
6	1x4	8	16	6 lb. 8d nails
2	1x6	10	10	
Rustic	1x6	10	40	
Rustic	1x6	8	200	
		Total	617	

TABLE V

"A" Type Portable Hog-House—Building Material for Plan "B" (Board and Battens).

Material as Bought			Material as Used		
No. of Pieces	Size of Stock (inches)	Length (feet)	No. of Pieces	Length (feet)	Use
2	4x6	10'	2	10'	Skids
5	4x4	8'	5	8'	Joist
1	4x4	6'	2	3'	Joist for runway
7	2x12	8'	7	8'	Flooring
1	2x6	8'	1	8'	Flooring
8	2x4	8'	8	8'	Plates and rafters
2	2x4	8'	4	3'9"	Purlins
1	2x4	10'	1	9'	Ridge pole
1	2x4	14'	1	2'6"	Door lintel
			2	2'	Window
			4	1'8"	Ventilator
2	2x4	10'	4	5'	Studs, window & door
1	2x4	10'	4	2'6"	End studs
2	1x6	10'	2	9'	Ridge boards
2	1x4	10'	6	2'4"	Door hurdle
			2	2'10"	
26	1x12	8'	18	8'	Roof
			8	Vary	Ends
25	1x4	8'			Battens
1	1x4	10'	2	3'2"	Window trim
			2	1'8"	
1	1x4	8'	2	2'2"	Window casing
			2	1'6"	
4	1x4	8'	4	7'8"	Corner trim
1	1x4	10'	2	3'2"	Door casing
			1	2'6"	
1	1x4	12'	2	3'9"	Door trim
			1	3'2"	
1	2x12	8'	3	2'6"	Runway floor

Summary				Miscellaneous
No. of Pieces	Size of Stock (inches)	Length (feet)	Board Feet	
2	4x6	10	40	4— $\frac{1}{2}$ "x10" bolts and washers 2—8" strap hinges 2—4" T hinges 3—3" gate hooks and eyes 2—4" gate hooks and eyes 3' wire chain 6'—1"x $\frac{3}{8}$ " strap iron 7 lb. 8d nails 4 lb. 20d nails
5	4x4	8	53	
1	4x4	6	8	
8	2x12	8	112	
1	2x6	8	8	
10	2x4	8	54	
4	2x4	10	27	
1	2x4	14	9	
2	1x6	10	10	
26	1x12	8	208	
1	1x4	12	4	
4	1x4	10	13	
30	1x4	8	80	
		Total	626	

LIBRARY
 SWINE HUSBANDRY
 IN IDAHO
 1920

TABLE VI
Bill of Materials for Shed Roof Field House

Material as Bought			Material as Used		
No. of Pieces	Size of Stock (inches)	Length (feet)	No. of Pieces	Length (feet)	Use
2	4x6	14'	2	14'	Skids
5	4x4	8'	5	8'	Joist
7	2x12	12'	7	12'	Flooring
1	2x6	12'	1	12'	Flooring
1	2x8	6'	2	2'8"	Window sill
7	2x4	12'	7	12'	Rafters
4	2x4	12'	4	12'	Plates
7	2x4	12'	7	7'2"	Studs, front
			7	4'2"	Studs, back
2	2x4	12'	2	6'	Studs, end
			2	5'3"	Studs, end
1	2x4	14'	1	6'10"	Studs, end
			1	4'6"	Door stud
			1	2"	Door stud
5	1x4	16'			Trim
1	2x4	8'	1	4'6"	Door stud
			1	3'6"	Door lintel
Summary					
No. of Pieces	Size of Stock (inches)	Length (feet)	Board Feet	Miscellaneous	
2	4x6	14	56		
5	4x4	8	53	10— $\frac{1}{2}$ "x12" bolts	
1	2x6	12	12	2—4-light sash, 8"x10" panes	
1	2x8	6	8	3 $\frac{1}{2}$ lb. 3d nails	
20	2x4	12	160	10 lb. 8d nails	
1	2x4	14	10	7 lb. 20d nails	
1	2x4	8	5		
5	1x4	16	27		
Sheeting			200		
Rustic V siding			300		
Total.....			\$31		
4 bundles shingles					

TABLE VII
Bill of Materials for Portable Loading Chute

Material as Bought			Material as Used		
No. of Pieces	Size of Stock (inches)	Length (feet)	No. of Pieces	Length (feet)	Use
12	1x4	12'	12	12'	Sides
2	1x6	14'	2	12'	Sides
			2	1'2"	Doors
2	2x4	12'	2	11'10"	Top
6	2x4	12'	12	3'4"	Sides, vertical
			6	5'6"	Joist
2	2x4	10'	10	2'	Braces
3	2x10	12'	3	12'	Floor
2	1x2	14'	10	2'6"	Cleats
1	2x10	12'	2	6'	Cart frame
1	4x6	8'	1	3'9"	Bolster
			1	3'	Prop
1	2x6	12'	2	1'6"	Lower door
			3	2'6"	Lower door
1	1x4	12'	4	3'	Doors
1	1x4	14'	12	1'2"	Doors
Summary					
No. of Pieces	Size of Stock (inches)	Length (feet)	Board Feet	Miscellaneous	
13	1x4	12	52		
2	1x6	14	14	6—4" strap hinges	
2	2x4	12	64	1—8" strap hinge	
2	2x4	10	13	14— $\frac{1}{2}$ "x6" bolts and washers	
4	2x10	12	80	16— $\frac{1}{2}$ "x4" bolts and washers	
2	1x2	14	Linear	2— $\frac{1}{2}$ "x14" bolts and washers	
1	4x6	8	16	10 lb. 8d nails	
1	2x6	12	12	2 lb. 16d nails	
1	1x4	14	5		
		Total	256		

TABLE VIII
Bill of Materials for Lifting Crate.

Material as Bought			Material as Used		
No. of Pieces	Size of Stock (inches)	Length (feet)	No. of Pieces	Length (feet)	Use
7	2x6	8'	3	8'	Sides and bottom
6	2x4	8'	4	7'8"	Sides and bottom
3	2x8	8'	6	7'8"	Sides
1	2x8	8'	3	8'	Sides and bottom
1	2x6	10'	4	2'6"	Doors and ends
1	2x4	10'	4	2'6"	Ends
2	2x4	14'	4	3'4"	Doors
2	2x4	10'	8	2'6"	Doors
3	2x6	12'	6	5'4"	Uprights and blocks
Summary					
No. of Pieces	Size of Stock (inches)	Length (feet)	Board Feet	Miscellaneous	
1	2x8	10	13	56"— $\frac{1}{2}$ "x1 $\frac{1}{2}$ " strap iron	
3	2x8	8	32	2 steel rings, 4" outside diameter, of	
3	2x6	12	36	$\frac{3}{4}$ " material	
1	2x6	10	10	7'4"x $\frac{3}{4}$ " iron rod	
7	2x6	8	48	4-6" strap hinges	
1	2x4	14	9	2-6" gate hooks	
2	2x4	10	14	16— $\frac{1}{2}$ "x4 $\frac{1}{2}$ " bolts and nuts	
6	2x4	8	32	88— $\frac{1}{2}$ "x4" bolts and nuts	
		Total	194	12— $\frac{1}{4}$ "x8" bolts and nuts	

TABLE IX
Bill of Materials for Castrating Crate.

Material as Bought			Material as Used		
No. of Pieces	Size of Stock (inches)	Length (feet)	No. of Pieces	Length (feet)	Use
1	2x8	10'	2	3'	End uprights
			1	4'	Top
1	2x6	4'	1	4'	Top
1	2x4	10'	4	3'2"	Legs
			2	3'2"	Cross braces
1	2x4	8'	4	1'9"	Leg braces
1	1x12	4'	1	3'9"	Shelf
Summary					
No. of Pieces	Size of Stock	Length	Board Feet		
1	2x8	10'	13 $\frac{2}{3}$		
1	2x6	4'	4		
2	2x4	10'	13 $\frac{2}{3}$		
1	2x4	8'	5 $\frac{1}{2}$		
1	1x12	4'	4		
		Total	39 $\frac{2}{3}$		

TABLE X
Bill of Materials for Hog Shipping Crates.

Material as Bought			Material as Used		
No. of Pieces	Size of Stock (inches)	Length (feet)	No. of Pieces	Length (feet)	Use
Crate for 100-pound hog: 1'2"x2'4"x3'10"					
1	1x6	12'	9	1'4"	Floor
1	1x6	10'	3	2'8"	Door
			1	1'6"	End
1	1x6	8'	2	3'10"	Sides at bottom
2	1x4	8'	6	2'7"	Sides
1	1x4	10'	6	1'6"	Top and ends
3	1x4	8"	6	3'10"	Sides
1	1x4	8'	4	1'4"	Top
			2	1'4"	Door cross pieces
1	2x4	8'	2	4'	Floor joist
Summary					
			1—2x4x8	6	Board Feet
			1—1x6x12	6	
			1—1x6x10	5	
			1—1x6x8	4	
			1—1x4x10	3	
			6—1x4x8	16	
				40	Board Feet
Crate for 250-pound hog: 1'6"x2'9"x4'4"					
2	1x6	8'	10	1'8"	Floor
1	1x6	12'	3	3'2"	Door
			1	1'10"	End
1	1x6	10'	2	4'4"	Sides at bottom
2	1x4	10'	6	3'	Sides (vertical)
1	1x4	12'	6	1'10"	Top and ends
2	1x4	14'	6	4'4"	Sides
1	1x4	10'	4	1'8"	Top
			2	1'8"	Door cross pieces
1	2x4	10'	2	4'6"	Floor joists
Summary					
			2—1x6x8	8	Board Feet
			1—1x6x12	6	
			1—1x6x10	5	
			3—1x4x10	10	
			1—1x4x12	4	
			2—1x4x14	9	
			1—2x4x10	7	
				49	Board Feet
Crate for 500-pound hog: 2'x3'4"x6'					
2	1x8	12'	10	2'2"	Floor
4*	1x4	12'	7	6'	Floor
1	1x6	14'	3	3'9"	Door
			1	2'4"	End
1	1x8	12'	2	6'	Sides at bottom
2	1x4	12'	6	3'7"	Sides, vertical
1	1x4	14'	6	2'4"	Top and ends
3	1x6	12'	6	6'	Sides
1	1x4	14'	4	2'2"	Top
			2	2'2"	Door cross pieces
1	2x4	12'	2	6'	Joists (floor)
Summary					
			5—1x6x12	30	Board Feet
			6—1x4x12	24	
			1—1x6x14	7	
			1—1x8x12	8	
			2—1x4x14	10	
			1—2x4x12	8	
				87	Board Feet

*The floor is made double thickness to support heavy hogs, the top layer of boards being laid lengthwise.

Feeding and Management

should not be used much for breeding before he is one year of age. When buying a boar, it is usually desirable to see him before the pig is at least five or six months old. It is better if a pig is selected at a younger age that he can be acclimated to the conditions if he is subjected to new surroundings and management. It is of advantage also to see the pig when the selection is made so the observations serve as guides in selection will be more de-

allowed to run with the sow herd. This applies to mature boars. Experimental evidence indicates bad results from such practice. Some boars that do settle often produce small litters and should not be run with the herd. A young boar should be used on one service a day for a short breeding season or on one every other day for a longer season. A boar should be allowed two services each day if the breeding season is long.

No sow need be bred more than once each year. In some cases, a young boar which was allowed to run with the herd previously farrowed an average of 8.7 pigs that varied from one to six pigs with an average litter. In another similar case a young boar which was allowed to run with the herd settled three sows out of 16 sows. The litters were small (*More detailed information on this subject is available at this Station.*)

A boar should be kept in a good sized lot which is equipped with a good shelter and quarters in winter. This lot should not be too small. The sows are kept. He should be kept on pasture during the winter.

Boars are irregular feeders. During the breeding season they are very easily. It is essential during this period to give them encouragement to stay on feed by keeping all day and by feeding, at regular hours, a combination of roughage and nourishing. The boar often refuses to eat the trough to sour. An ample amount of clean water should be available. Fighting boars are often difficult to handle. They should be quiet if a barrow or a bred sow is kept with them.

There should be a definite grain combination for all boars. The ration should be varied to suit local peculiarities. The ration should be varied to suit local peculiarities. Changes in the feed should be made slowly. A sudden change throws the boar off feed. The boar should not be kept on a single ration. Either condition may impair his breeding. If a young boar is on pasture, he should be fed a ration of concentrates to cause him to grow normally. The maintenance ration should be about one to two pounds of concentrates each day. The boar should be about one to two pounds of concentrates each day. The boar should be about one to two pounds of concentrates each day when he is not in service.

by Weight
95
5

service
changes
ions are
al:

by Weight
20
5
5
5
3

he rate
Corn is
e some
in the
rt.

immed
ng sea-
he feet
ey will

y is to
ness is
gain-
repro-

cality,
nprise
cussed
proper
intain
ixture
orage.
grain.
more
than

and is in pasture. He should be fed enough so he will not lose weight in the breeding season. In fact, he should be gaining a little weight. A variety of home-grown grains, with skim milk, shorts or tankage and green feed, if available, will give good results during the breeding season. One-half pound of salt should be added to each 100 pounds of feed.

When the boar is not in service he may be fed rather liberally.

Ration 1

	Parts by Weight
Barley, wheat or corn.....	1
Skim milk	1
Forage or alfalfa hay	

Ration 2

	Parts by Weight
Barley, wheat or corn.....	1
Tankage	1
Forage or alfalfa hay	

It is almost impossible to make up a ration for boars in the winter that will prove adequate for the entire season, as it seems that different rations are often necessary to keep them on feed. The following rations are suggested because they have been found satisfactory in general.

Ration 1

	Parts by Weight
Barley	1
Wheat	1
Oats	1
Skim milk	6

Ration 2

	Parts by Weight
Barley	1
Oats	1
Shorts	1
Corn (yellow)	1
Tankage	1

Skim milk may be substituted for tankage in ration 2 at the rate of 12-15 pounds of skim milk for each pound of tankage. Tankage is very palatable to swine. It is therefore often desirable to have it available to use in boar rations. In corn producing sections state corn may be substituted for the cereals, at least in part.

The feet of boars, especially heavy boars, should be trimmed back so they will not break off as they often do in the breeding season. Some boars are useless until such injuries are healed. They should not be trimmed so short as to expose the "quick" or the hoof, or become tender.

Sows. One of the essentials to success with the brood sow is to keep her in a vigorous condition of health. The degree of fatness is not an assurance of vigor. When the sow is bred she should be in good condition. This has a practical effect in stimulating her reproductive functions.

The kind and amount of grain to feed will vary with the age and condition of the sows. Home-grown feeds should constitute the bulk of all rations. Any one of the grains previously discussed that is produced most economically may be used with the proper supplements. Mature, dry sows in good condition will maintain their weight on one to one and one-fourth pounds of a grain mixture per day for each 100 pounds of live weight when they are on feed. Thin mature sows often need more than this amount of feed. Growing gilts intended for breeding purposes should be fed liberally, but not all they will eat. When they get a little less

they will eat they ordinarily take more exercise and stay in a more thrifty condition. Well grown gilts may be bred when eight months of age.

During the winter gestation period, the brood sow's ration is important. Very little forage is available during this period, hence a substitute for it should be provided whenever possible. A high quality alfalfa hay, although not equal to forage, has been successfully used. Alfalfa hay may be fed in racks to good advantage (Fig. 16). This method stimulates the sows to take more exercise and helps them keep in good condition. Alfalfa hay is high in minerals and vitamins. Sows getting alfalfa hay will do well on a variety of grain combinations. Wheat, barley, or corn alone or in combination, if available, may serve as the basis for this ration. Any of the grains may be replaced in part by oats for pregnant sows, and in pea producing areas one-third of the grain may be replaced with pea screenings. Too much emphasis cannot be placed on the advisability of feeding alfalfa to pregnant sows, and oats is always a good feed for open, bred, or suckling sows.

Suggested Feed Combinations for Wintering Sows

Ration 1

	Parts by Weight
*Barley	1
**Skimilk	1
Fine alfalfa hay in rack	

Ration 3

	Parts by Weight
Barley	1
Wheat	1
Oats	1
Skimilk	3
Fine alfalfa hay in rack	

Ration 5

	Parts by Weight
Barley	8
Peas	2
Skimilk	6
Fine alfalfa hay in rack	

Ration 2

	Parts by Weight
Barley	16
Oats	6
†Tankage	1
Fine alfalfa hay in rack	

Ration 4

Barley	16
Oats	8
Shorts	4
Tankage	1
Fine alfalfa hay in rack	

Ration 6

	Parts by Weight
Barley	16
Oats	8
Peas	3
Tankage	1
Fine alfalfa hay in rack	

Time to Farrow. Farmers have more time available to devote to the care of pigs when they are farrowed before the spring work. Early-farrowed pigs can utilize hogging-off crops, such as peas, in time to reach the early market. Early farrowing also offers an opportunity to produce two litters per year whenever that is deemed

* Wheat or corn may obviously be substituted for barley.

**When tankage is used to replace skimilk figure one pound of tankage (60 per cent) equivalent to 12 to 15 pounds of skimilk.

† Fish meal may be substituted in equal amounts for tankage, and if linseed oil meal is used as part of the supplement, two pounds should be allowed for each pound of tankage replaced.

advisable. This method requires, as a rule, more expensive and warmer houses. Pigs farrowed early, however, are subject to a nutritional disease called "Anemia" (See this subject under "Swine Diseases" for full discussion).

Care at Farrowing Time. The lack of proper sanitary measures is responsible for a large percentage of pig losses. Most rigid sanitary measures are necessary to destroy the little pig's worst enemy, the round worm. It has been found that round worms are very prevalent, especially in herds which have been kept in the same lots for a number of years. Many pigs die very young from round worms (See "Swine Sanitation" for fuller discussion of preventive measures).




Fig. 16.—Hay rack. A practical way to feed alfalfa hay. Note cover on rack. When hay becomes wet, the flavor of the hay is largely destroyed. Hay should be fed in rather small quantities and renewed at short intervals. Sows spend much time around these racks in all kinds of weather getting valuable exercise as well as very necessary food constituents.

Sows that are accustomed to exercise may become constipated if left in a small pen where there is little opportunity for exercise. The feeds should be made laxative by adding some bran. The amount of feed may also be reduced, but not to the point where the sow becomes unduly hungry. Constipation generally sets up a feverish condition which often results in viciousness, and in many cases pig eating. Pig eating may also occur if the sow has not been fed enough minerals or protein during the pregnancy period. It is not natural for a sow to eat her pigs. Constipation, feeding grain alone, not enough protein or minerals, and allowing her to eat her afterbirth or laidover pigs often cause her to develop the habit (See Fig. 17 for farrowing rails).

TABLE XI

Gestation Table.

The following table is based on a 112 days gestation period. If, for instance, a sow is bred February 1, she is due to farrow May 24. If she is bred May 24, she is due to farrow September 13. The date to the immediate right of the date on which she was bred is her date to farrow. Breeding a sow November 9 to 11 will usually bring March 1 pigs.

Feb.	May	Sept.	Jan.	Apr.	Aug.	Dec.	Mar.	July	Nov.	Feb.	June	Oct.	Jan.
1	24	13	3	25	15	5	27	17	6	26	18	8	28
2	25	14	4	26	16	6	28	18	7	27	19	9	29
3	26	15	5	27	17	7	29	19	8	28	20	10	30
4	27	16	6	28	18	8	30	20	9	Mar. 1	21	11	31 Feb. 1
5	28	17	7	29	19	9	31 Apr. 1	21	10	2	22	12	2
6	29	18	8	30 May 1	20	10	2	22	11	3	23	13	3
7	30	19	9	1	21	11	2	23	12	4	24	14	4
8	31 June	20	10	2	22	12	3	24	13	5	25	15	5
9	1	21	11	3	23	13	4	25	14	6	26	16	6
10	2	22	12	4	24	14	5	26	15	7	27	17	7
11	3	23	13	5	25	15	6	27	16	8	28	18	8
12	4	24	14	6	26	16	7	28	17	9	29	19	9
13	5	25	15	7	27	17	8	29	18	10	30 July	20	10
14	6	26	16	8	28	18	9	30	19	11	1	21	11
15	7	27	17	9	29	19	10	31 Aug. 1	20	12	2	22	12
16	8	28	18	10	30	20	11	1	21	13	3	23	13
17	9	29	19	11	31 Sept. 1	21	12	2	22	14	4	24	14
18	10	30 Oct.	20	12	2	22	13	3	23	15	5	25	15
19	11	1	21	13	2	23	14	4	24	16	6	26	16
20	12	2	22	14	3	24	15	5	25	17	7	27	17
21	13	3	23	15	4	25	16	6	26	18	8	28	18
22	14	4	24	16	5	26	17	7	27	19	9	29	19
23	15	5	25	17	6	27	18	8	29	20	10	30	20
24	16	6	26	18	7	28	19	9	29	21	11	31 Nov.	21
25	17	7	27	19	8	29	20	10	30 Dec.	22	12	1	22
26	18	8	28	20	9	30	21	11	1	23	13	2	23
27	19	9	29	21	10	31 Jan.	22	12	2	24	14	3	24
28	20	10	30	22	11	1	23	13	3	25	15	4	25
.....	21	11	31	23	12	2	24	14	4	26	16	5	26

During the process of farrowing, an attendant should be at hand, especially if farrowing takes place in cold weather. If pigs chill at the time of farrowing, their chances for a good start have been reduced. A basket, barrel, box, or other suitable container should be at hand in which warm bricks covered with straw, or in which a jug containing warm water has been placed (*Fig. 18*). The pigs should be rubbed gently with a burlap bag before being placed in the container. The mucous should be removed from the mouth.

The pigs may be taken to the sow for nursing when they are dried off, after which they may be returned every two hours until they have gained enough strength and acquired sufficient control to get along without additional assistance. Sows that are too vicious to allow the application of this detailed management may be profitably replaced by others that are gentle.

A pig is born with eight small but very pointed tusk-like teeth, two on either side of the upper and lower jaw immediately back of




Fig 17.—Farrowing rails are valuable. They are, as a rule, 10 inches from the floor and eight inches from the wall.

the front teeth. It is generally considered advisable to clip these teeth. This can be done with a pair of side-cutting nippers (*Fig. 19*). These teeth should not be pulled out nor imperfectly cut off. Sharp projections left on the teeth often cause considerable annoyance in cutting the lips and tongues of young pigs and the teats of the sow. The jaw or gum should not be injured when cutting the "needle teeth." Make a smooth cut and do not draw blood. Cuts in the jaw or gums and tongue are avenues of invasion for various types of infection that often cause sore mouth.



Fig. 18.—In cool weather pigs may be kept comfortable in a box until they are strong enough to be left with the sow.

Marking the Pigs

It is necessary to mark purebred pigs so they will be properly identified if they are to be recorded. Since metal tags are easily pulled out of the ears, it is more practical to use one of the well recognized systems of ear notching. Notches may be cut into the ear with a knife when the pig is small. It is preferable, however, to use a special ear-notcher (*Fig. 19*). Pigs should have their ears notched before the litters are allowed to intermingle. A good rule to follow is to mark the pigs the first or second day. In *Fig. 20* will be found a satisfactory method for marking.




Fig. 19.—A. Ear marker for notching the ears. B. Nippers commonly used for cutting "needle teeth" in young pigs.

Care of the Sow and Pigs

The sow should not be urged to eat during the first 24 hours after farrowing. Some sows crave feed sooner. The feeding should be rather light the first few days. Provide clean water, however, preferably with the chill removed. There is no set rule that applies to all sows during this period of management. The attendant must study the habits of each sow. The object should be to feed enough so the sows will not become unduly hungry, and yet not so much that the flow of milk will be stimulated too much before the pigs have developed a little size and are capable of using it all. Too much milk the first few days often develops scours in the little pigs. Scours may also be caused by filth, drafts, chills, damp bedding, and often by changing the sow's ration suddenly. The cause for scours




Fig. 20.—(At left) Value of notches in right and left ears. (At right) Ears notched to indicate No. 73.

	Left Ear	Right Ear
Lower side, next to head	1	10
Lower side, midway from tip to base of ear	3	30
Lower side, tip of ear	5	50
Upper side, next to head	2	20
Upper side, midway from tip to base of ear	4	40
Upper side, tip of ear	6	60

should be removed first (*For specific treatment see "Scours" under "Swine Diseases"*). One-half of the sow's ration during the period she is nursing the pigs may contain shorts or middlings, which seem to stimulate the milk flow. The feed may be made up into a slop to encourage the consumption of water. The allowance should be cleaned up at each feeding because remaining residues may become stale and sour.

Results from work done at this Station on milk production in sows indicate considerable variation in the amount of milk that sows produce. Some sows produce almost two gallons in twenty-four hours when in full milk. Sows with such heavy production obviously will require considerable feed. Other sows produce scarcely one gallon of milk each day.

The pasture should be on clean ground where hogs were not kept the previous year. The individual house and troughs in the pasture lot should be scrubbed with boiling water (*See Swine Sanitation*). The pigs and sow should be crated and moved to the new pasture. If the sows are driven, they may pick up worm eggs by dragging the udder over contaminated ground. Little pigs should not be kept on the same ground two years in succession, nor should they be kept on ground which has been occupied by hogs the previous year. This plan of management requires more pasture lots than are required when the same lots are utilized during successive years. In alternate years, however, the pasture lots may be cut for hay. The total expense may be somewhat higher, but years of carefully conducted experimental work show clearly the practicability of this program.

Rations for the Sow. Sows lose as a rule, about three-fourths of a pound a day when they are suckling pigs. Heavy milkers will lose more. When the pigs are from ten days to two weeks old, the sow should be on full feed and be given all she will clean up. If she loses weight too rapidly, feed her three times a day. Any one of the following rations may be used:

**Ration 1*

	Parts by Weight
**Barley	100
Shorts	25
† Tankage	4

Ration 2

	Parts by Weight
Barley	100
Peas	20
Tankage	4

Ration 3

	Parts by Weight
Barley	100
Wheat	100
Tankage	12

Ration 4

	Parts by Weight
Barley	100
Oats	50
Shorts	25
Tankage	8
Or skimmilk	150





Fig. 21.—(Upper) It is advisable to feed nursing pigs in a creep. (Right) Adjustable openings for creeps are in common use. They can be adjusted as the pigs develop.

* Before the sow gets out on pasture add alfalfa leaves to the extent of 5% to any ration selected.

**Barley, corn or wheat may be used.

† When skimmilk is available use 12 to 15 pounds in place of each pound of tankage or fish meal.

Creep Feeding. Creeps should be made available for the pigs by the time they are three weeks old (*Fig. 21*). At this time it is preferable to give them dry feed in the form of coarsely ground wheat, or a combination of coarsely ground wheat and cracked corn. After they have become accustomed to eating, the feed may be changed to a mixture of ground wheat, corn or barley, shorts or middlings, ground oats, skim milk or tankage fed in the form of slop. The troughs must be watched so feed is not left to become sour. A good ration may be made up of ground corn, barley, or wheat, 70 parts; shorts 26 parts; and digester tankage, 4 parts. The protein supplement may be fed entirely in the form of dairy by-products by allowing 1 to 2 pounds of skim milk for each pound of grain mixture. This ration will be suitable for the weaning age also. If the price of oats is comparable with the price of other grains, it may be fed to replace one-half of other grains.

Weaning. At eight to ten weeks of age the pigs are ready to be weaned. The sow should be removed from the pigs, leaving them undisturbed in the lot with which they are familiar. The sow's ration should be reduced so the milk flow will be discouraged. Whole oats may be fed to the sow to advantage at this time. If the milk flow persists and the udder becomes caked, it should be bathed with warm water, dried off, and massaged with equal parts of lard and turpentine. Her management during the summer is comparatively simple. If she has been suckled down she should be fed grain in addition to the forage until she gets back to a thrifty condition, after which she will not need much grain so long as the forage is good.

Castration. The boar pigs should be castrated preferably a week or ten days before they are weaned. Pigs that are castrated and weaned at the same time usually undergo too great a shock. Castration should always be done under sanitary conditions and a three per cent solution of a standard coal tar livestock dip should be used for the scrotum and the wound. The cut should be made low so as to allow efficient drainage (*See Fig. 13 and Table IX*).

Developing the Weanling Pigs. The method used in developing the pigs after weaning will have to be governed largely by the time at which they are expected to be ready for market. They must be given a full grain ration if they are to be marketed at the earliest possible date. Lighter grain feeding may be desirable if they are to be developed as feeders, with a view of hogging off crops, or cleaning stubble fields. The bulk of the grain which the pig that is crowded for the early market consumes has been carried over from the previous year. The pig which is developed for use in the stubble field will consume much more of the new crop. Generally speaking, the areas in the state that are suitable for the production of feeder pigs are also suitable for finishing them. Very few feeder pigs are sent out of Idaho.

Feeding the Market Hog on Forage. Pigs may be carried along on forage with a limited amount of grain or they may be fed a full ration with a view of preparing them for the early market. The

amount to feed will depend upon the date one has in mind to market the hogs. Pigs on good forage, when allowed all the grain they will consume, will usually gain about 1.25 pounds per day, making it possible for early March pigs to reach a satisfactory market weight by the first of September. Four pounds of grain a day for each 100 pounds of weight in hogs is considered a full ration. Thus a 50-pound pig on a full ration will consume about two pounds of grain a day. Pigs fed a three-fourths ration, or three pounds of grain for each 100 pounds of weight will gain about one pound per day on forage. They will gain three-fourths pound per day if they receive two pounds of grain for each 100 pounds of live weight, and if fed only one pound for each 100 pounds of weight, they will gain only one-half pound per day. A number of factors determine what method to use, such as the amount of feed available, the comparative cost of feed, and the price of market hogs, the time of market, the availability of waste feed to be harvested by the hogs, and the cost of carrying pigs. A choice of limited rations may be desirable if the carrying cost of pigs is small and one is not interested in early markets.

One lot of March pigs at this Station, fed a full grain ration while on forage, went to market eleven days earlier and sold for \$1.25 per hundred weight more than did a similar lot which was fed a grain ration limited to about three-fourths of what they would eat. The lot on a full ration cost slightly more but the increased sale price off-set this by more than \$1.50 per pig in favor of the full-fed lot.

Rations made up of a variety of grains are, as a rule, preferable to any of the grains alone and to these should be added one or more protein supplements. The following are suitable combinations for full-fed pigs on forage:

Ration 1

	Pounds
Barley	75
Oats	25
*Tankage	4

Ration 2

	Pounds
Barley	75
Shorts	25
Tankage	3

Ration 3

	Pounds
Barley	30
Wheat	30
Oats	20
Shorts	20
Tankage	2

Ration 4

	Pounds
Barley	30
Wheat	30
Peas	20
Oats	20
Tankage	2

The most economical gains on the basis of feed requirements for pounds of grain, are made when the pigs are young. Table XII will give some information on the daily consumption of feed and requirements for 100 pounds of gain.

Dry Lot Feeding. Usually the hogs have to be finished for market in the dry lot. With pigs that have been on full feed during the

*If skimmilk is used in place of tankage, use 15 pounds of skimmilk for each pound of tankage replaced.

TABLE XII
Relation of Weight of Pigs to Feed Consumed and Rate of Gain.*

Wt. of pigs	Actual	Average feed	Feed eaten	Average	Feed for 100
	average wt.	eaten per day	daily per 100 lbs. live wt.	daily gain	pounds gain
lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
15-50	38	2.2	6.0	.8	293
50-100	78	3.4	4.3	.8	400
100-150	128	4.8	3.8	1.1	437
150-200	174	5.9	3.5	1.2	482
200-250	226	6.6	2.9	1.3	498
250-300	271	7.4	2.7	1.5	511
300-350	320	7.5	2.4	1.4	535

*"Feeds and Feeding"—Henry and Morrison.

forage season, the period for dry-lot feeding need not be long, and the following rather simple combinations may be used:

Ration 1

	Pounds
Barley or wheat	100
Tankage	8
or skimmilk, 1 pound for each pound of grain.	

Ration 2

	Pounds
Barley or wheat	90 to 95
*Oats	5 to 10
Tankage	6 to 8
Or skimmilk as in 1.	

*It is likely that in some cases the oats may make the ration bulky and not fattening enough for the growing hogs. If this is the case, reduce the amount of oats.

In the event the pigs have not been getting a full grain ration on forage and a longer dry lot feeding period is necessary, at least 5 per cent of the ration should be alfalfa leaves or meal.

TABLE XIII
Amount of Different Protein Supplements Required to Balance Barley for Pigs on Legume Pasture.*

Protein Supplement	Parts by Weight		
	Barley	Supplement	Nutritive Ratio
Skimmilk (Centrifugal)	100	100	1:6.03
Buttermilk	100	100	1:6.09
Tankage (60% protein)	100	5	1:6.03
Standard wheat shorts	100	50	1:6.26
Linseed-oil Meal (o. p.)	100	12	1:6.04

*Calculations based upon digestible nutrients—Henry and Morrison's "Feeds and Feeding." Approximately the same nutritive ratio will obtain if wheat is substituted for barley.

Developing Pigs for the Breeding Herd. One should make the best selections possible when gilts are retained for the breeding herd. The selection should be made before the pigs are placed on a full finishing ration. A high degree of finish in gilts is harmful and unnecessarily expensive. Gilts intended for breeding purposes should always be allowed access to a good forage crop and should be fed a sufficient amount of grain to keep them in thrifty condition. Oats should be used liberally for growing pigs that are developed for breeding purposes. The grain ration used should be supplemented with skimmilk or tankage. Gilts should be developed to weigh from 200 to 250 pounds when they are bred. They should not be bred until they are eight months old so they will farrow when they are one year old.

Swine Diseases, Prevention and Control

Swine production is one of the established livestock enterprises of Idaho. The long continued production of, or the rapid increase of livestock in a locality is usually accompanied by more or less trouble with diseases. The farmer in Idaho is aware of the fact that he can make a profitable enterprise of swine production providing hog diseases do not interfere with his operations. He is, therefore, interested in knowing what the most common and most destructive swine diseases are; how they are introduced onto his farm; how he can prevent their entrance; how he can eliminate or hold in check diseases already present on the farm; and what he can do in the way of treatment for disease conditions present in his hogs.

Prevalence of Swine Diseases in Idaho

The acute infectious diseases of hogs are not generally prevalent in Idaho. Hog cholera, hog flu and hemorrhagic septicemia occur only occasionally. Tuberculosis of the avian or fowl type is, however, quite common in practically all sections of the state.

The intestinal parasites and filth-borne diseases of swine are responsible for the greater portion of losses from diseases in this state. Owing to the nature of their attack they are too often overlooked. Intestinal round worms, lung worms, coccidiosis, naval infection, necrophorus infection and other soil-borne infections associated with enteritis and diarrhea of pigs comprise the common sources of trouble.

Hog lice are very common, while hog mange is rare. The common nutritional diseases of swine such as rickets, goitre (hairlessness) and anemia also occur.

Swine Sanitation

The one outstanding weapon that can be economically and satisfactorily used in combating swine diseases is cleanliness in raising the pig crop.

The average swine producer is aware of the fact that little trouble is experienced in growing disease-free hogs during the first year of a swine-producing enterprise. As the grower improves the type of his swine and gains in knowledge in feeding and management during the first few years of production, he expects better pigs. However, the number of runty pigs and the amount of disease, not accounted for by hog cholera or other specific acute infections, usually increases. Either he becomes discouraged and goes out of the business, or gets additional information regarding the causes behind his failure to grow pigs successfully. Factors responsible for the loss of baby pigs and the development of runts are numerous but can be explained on a very simple basis and can be controlled efficiently by the swine producer.

During the years between 1915 and 1925, the great hog producing areas of the Middle West were finding it practically impossible to successfully raise even fifty per cent of the pigs farrowed. On many farms, hog production was discontinued.

In 1919 under the direction of Dr. B. H. Ransom of the United States Bureau of Animal Industry, an investigation of the cause of this trouble was begun. It was found, in the first place, that most growers having such trouble were yarding their pigs in permanent hog lots. It was also found that many of the affected pigs had pneumonia or thumps and upon a post-mortem examination many immature round worms were found in the lungs. Sore mouths (stomatitis), lung worms (those worms that mature in the lung), necrotic enteritis ("necro"), and other disease conditions of a comparable nature were found associated with this round worm infestation in the lungs and intestines of pigs grown in permanent hog lots.

These findings led to the establishment of a hog-lot sanitary program participated in by a large number of swine producers. The results obtained by the use of this sanitary system was so satisfactory that it has been generally adopted. This sanitary system is very completely outlined in Leaflet No. 5 of the Bureau of Animal Industry dealing with "The Prevention of Round Worms in Pigs." It emphasizes the following steps:

First, thoroughly clean the farrowing pen. Remove all loose litter and equipment and wash the house out with boiling water to which has been added one pound of lye to 20 gallons of water. Equipment should be washed in a similar manner.

Second, wash the sow off with warm soapy water, removing all dirt from the body and feet and place her in the clean pen.

Third, keep the pigs confined to the farrowing pen if it is in the vicinity of old hog lots, until the sow and pigs are moved to a clean pasture.

Fourth, haul the sow and pigs to pasture to avoid picking up worm eggs from contaminated soil.

Fifth, prevent the pigs coming in contact with contaminated ground by the use of hog-tight fences.

Sixth, do not run pigs on low wet pastures or allow them on a pasture that is being irrigated at the time.

Seventh, prevent the development of polluted areas near the feeding equipment by moving such equipment occasionally or by using feeding platforms. (Fig. 22).

This hog lot sanitary system is aimed primarily at the prevention of round worms but successfully controls thump or pneumonia, bull nose, enteritis, and other "necro" infections, lung worms, kidney worms, coccidiosis, and other filth-borne infections of hogs. It also assists materially in preventing the development of hog cholera, flu, and other acute infectious diseases.

Advantages of Hog Lot Sanitation

In order that growers be fully aware of the importance of sanitation in swine production, the following data from University of Illinois Circular No. 306 are presented:

This information is based on records of 13,478 pigs raised from 1,977 sows on 154 farms in Illinois by the use of the sanitary system and compared with a large number of pigs raised in permanent hog

lots on the same farms. Ninety-eight and two-tenths per cent of the pigs saved at farrowing time were raised by use of the sanitary method.

Large Litters Raised. An average of 1.6 to 2.7 more pigs per litter were raised. Farms formerly having the greatest trouble increased the size of litters most.

Number of Sows Needed Reduced. Twenty-eight per cent fewer sows were needed to produce the same number of pigs. Seven or eight sows under proper sanitary conditions thus raised as many pigs as ten sows with poor sanitation.




Fig. 22.—The sanitary features of the feeding platform are indispensable if hogs are to be fed for a protracted period in one place.

Almost No Runts with Sanitation. In addition to the number of pigs saved by sanitation, the number of runts was reduced from 18 out of every 100 pigs to one out of 100 pigs. A runt pig is usually a liability.

Faster Gains by Sanitary System Pigs. The sanitary system pigs weighed an average of 27 pounds more at four months of age than permanent hog lot pigs. Furthermore, sanitary system pigs reached marketable size seven weeks earlier and weighed twenty-eight pounds more than the permanent-hog-lot pigs.

Preventing the Introduction of Diseases

Hog diseases do not develop spontaneously. Any hog disease in order to develop in a herd must first be introduced just as any weed in the garden or field must be started by first introducing some weed seed. Some swine diseases, as hog cholera, develop rapidly after introduction. Others, as "necro" and intestinal parasites, develop slowly and may not cause great damage until they have been allowed to develop for two or three years.

Most hog diseases are introduced in the bodies of carrier animals;

that is, animals that have the disease. Consequently, the introduction of affected animals from outside sources is the most common cause of new diseases in a herd.

Recently, a large number of hogs have been shipped into Idaho both as breeding stock and feeders. Such procedure is a very serious menace to the swine producing industry of Idaho. Many of these hogs have died of disease, and unknown and unsuspected numbers of parasites and disease organisms have been introduced that will continue to afflict our herds. One man shipped in 89 hogs of which 45 died during the first month. In addition, 80 of 150 shoats already on the farm died of diseases introduced with this shipment. Round worms, lung worms, coccidiosis, and other parasites may also have been introduced. These diseases make it more difficult to grow hogs there in future years.

Federal and state sanitary officials prohibit the inter-state movement of noticeably diseased animals, but they cannot be sure that many chronic diseases such as round worm, "necro," and coccidiosis are not being carried in the bodies of healthy appearing animals. Canada prohibits the importation of hogs that have been vaccinated with hog cholera virus or that come from an area affected with cholera. Development of hog cholera has thus been successfully prevented. Idaho requires that all hogs from a section where hog cholera is prevalent must be vaccinated twenty-one days before they are permitted to enter this state. They must also be dipped.

The swine grower should supplement these precautions by being extremely careful in regard to the source of additions to the herd even though they come from a neighboring farm. Hogs from such a farm may harbor the organisms responsible for "necro," round worms, or even tuberculosis without showing evidence of it. If possible, only boar pigs should be introduced into a herd from outside sources, and they should be isolated from the herd for some time before being placed in contact with it.

Some diseases such as tuberculosis and bang abortion disease may be contracted from other species of livestock on the farm. Avian or fowl tuberculosis is a very common disease of swine in Idaho and is contracted from tubercular poultry. Bang abortion disease may be contracted from infected cattle as may also the bovine form of tuberculosis.

Elimination of Diseases in the Herd

Every precaution should be taken to prevent the introduction of swine diseases into the herd, but much may be done in eliminating diseases already present in the herd. Animals visibly sick should be immediately segregated from the remainder of the herd to prevent the disease spreading, in case it is of a contagious nature. Most infectious diseases as well as practically all parasitic disturbances of swine are transmissible, that is, "catching." The early isolation and treatment of animals affected with cholera, flu, hemorrhagic septicemia, mange, scours, or "necro" will do much to prevent the disease spreading to other animals in the herd. The inauguration of a swine

management plan that provides for cleanliness in the young growing pigs will do much to prevent the development of intestinal parasites and filth-borne diseases commonly present in Idaho swine herds.

The swine producer is not capable of detecting all animals affected with intestinal parasites or filth-borne diseases. Since these diseases are of major importance to the swine grower of Idaho, a definite program must be followed to prevent their spreading to the young stock. These diseases are preventable but are in general not amenable to treatment.

Control of Hog Diseases

Diagnosis. Diagnosis or the determination of the exact disease affecting hogs is of primary importance in deciding upon the line of treatment to be used. It is, therefore, highly desirable that the local veterinarian be called into consultation early. The opinion of other agencies such as the Federal or State Livestock Officials may be obtained. Your county agent will be able to assist you in getting in touch with these sources of information. Considerable information concerning the diagnosis and treatment of hog diseases may be obtained from Farmers' Bulletin No. 1244 on "Diseases, Ailments, and Abnormal Conditions of Swine," which may be obtained from the extension service of the Federal Government or from the University Extension Division, Boise and Moscow, Idaho.

Diseases such as cholera, hemorrhagic septicemia, tuberculosis, abortion, and anthrax are difficult or impossible to treat. Other diseases such as "necro," round worms, lung worms, and coccidiosis, pneumonia, poisoning, rheumatism, and filth-borne diseases are best controlled by following a rearing and management program that will prevent their development.

The swine producer should never place great faith or much confidence in treatment as a means of controlling intestinal parasites. No safe and satisfactory treatment has been developed for the control of lung worms of hogs. Furthermore, no medicine has been found to be satisfactory in the prevention and control of coccidiosis of hogs. Coccidiosis is a very common disease of young hogs in practically every hog producing locality.

Recently a pig from a herd of 200 shoats that were not doing well was received at the laboratory and on autopsy was found to harbor intestinal round worms and lung worms. It was also affected with coccidiosis. Hog-lot sanitation is the only practical solution of these conditions.

Hog Cholera. Hog cholera is an infectious herd disease of hogs. Its symptoms are high fever, loss of appetite, excessive thirst, and the development of a typical hog cholera squeal. If cholera is suspected, a veterinarian should be called immediately since the only known method of control is through vaccination. All pigs in contaminated areas should be vaccinated with the double treatment soon after they are weaned. Vaccination by the double treatment confers life immunity against cholera.

Hog cholera can be prevented from gaining entrance to a herd by guarding against the three main channels through which it is introduced. It is most often introduced by adding breeding stock or feeders from outside sources. About 40 per cent of the outbreaks are estimated to be initiated through the injudicious use of hog cholera virus, particularly through farmer vaccination. The third common source of infection is through pork products in garbage fed to hogs. By guarding against these main avenues of entry, hog cholera can be effectively prevented.

Medical treatment is not efficient in the treatment or control of this disease. It may be prevented by vaccination or by preventing its introduction (*See Farmers' Bulletin No. 834, on Hog Cholera*). Vaccination for hog cholera is most effective on healthy normal pigs. Swine affected with worms or other debilitating diseases often suffer heavy losses from hog cholera vaccination. Many veterinarians prefer to vaccinate some hogs with a mixed infection vaccine at the time of hog cholera vaccination.

Tuberculosis. The avian or fowl type of tuberculosis of hogs is a very mild form of the disease, most of the lesions being located in the glands of the head. The absence of bovine or cattle type of tuberculosis in hogs in this state is accounted for by the absence of this disease in cattle except in an occasional herd. Recently, at a local United States inspected packing plant the heads of 33 out of 39 hogs were condemned for human food because of infection with avian tuberculosis. In general, the farmer stands such a loss indirectly, if not directly, for no packing plant can afford such losses unless it pays less for live hogs. Hogs can be prevented from contracting avian tuberculosis by eliminating all affected chickens from the farm (*See Farmers' Bulletin No. 1200 on Tuberculosis of Fowls*).

Hog Flu. Hog flu is a recently discovered herd disease of swine. All animals in the herd come down with the disease within a few days and become extremely sick, refusing all feed. About the time that one expects them to begin dying they make a turn for the better and all but from two to five per cent recover. It is at first often confused with hog cholera, but the low mortality soon corrects this error.

The use of one ounce of magnesium sulphate (Epsom salts) for each 100 pounds of live weight as a laxative, good nursing and an alkaline mixture in the feed seems the only satisfactory treatment. The addition of one ounce of a mixture of equal parts of sodium hydroxide and sodium bicarbonate to each gallon of water in which the whole grains to be used are soaked and fed once daily has proved a satisfactory alkalinizing procedure.

Necrophorus Infection. Necrophorus infection or "necro" and filth-borne diseases are prevalent wherever hogs have been raised extensively without following proper sanitary precautions. There is some question as to whether or not bacillus necrophorus is the initial cause of all these afflictions, but it is quite definitely known that it at least plays a secondary role in their cause. This organism

as well as those organisms responsible for colds and pneumonia and navel infections in young pigs, pus infections of the glands of the head and neck, and joint infections in older pigs are the cause of extensive losses to hog growers where the hog lots are used continuously year after year. No satisfactory treatment has been discovered for these conditions. However, they are satisfactorily and easily prevented by following a sanitary program.

A recent communication from a farmer in the Camas Prairie section of northern Idaho states that of two litters of spring pigs all but three died of bull nose ("necro" of the head) and that one of three litters of three weeks old fall pigs was then affected.

Navel infections that ultimately result in arthritis or joint infection as late as several weeks after birth are often fatal.

A large percentage of hogs slaughtered at the local United States inspected packing plant are affected with abscessed glands or joint infection causing a great loss of otherwise valuable meat. Many of such infected hogs die and others never become sufficiently finished to be sold for meat. These losses, though not easily seen because of the chronic nature of the diseases causing them, in the long run represent one of the major leaks in the profits of the swine producer. Since these conditions do not respond to treatment, the logical procedure is to prevent them from developing by following a sanitary method of rearing (*See Swine Sanitation*).

Scours, Diarrhea and Enteritis. Diarrhea or scours is usually caused by inflammation of the intestines—a condition known as enteritis. Scours may develop in baby pigs as a result of faulty feeding of the sow (*See Feeding and Management*). However, filthy and unclean farrowing pens or mud and filth on the udder of the sow are responsible for most of these cases.

Older pigs may develop bloody diarrhea or scours from an infection with coccidiosis, necrotic enteritis or worm infestations. Occasionally scours may be brought on by the pigs eating spoiled feed, poisonous plants, lye, white lead or other noxious material that may be taken in the feed.

Most of these conditions may be prevented by following the sanitary program previously outlined and by carefully guarding the source of feed.

Most of these disease conditions are not amenable to treatment. However, some of the symptoms may be relieved by the judicious use of medication. Epsom salts at the rate of one ounce for each 100 pounds of pig or the use of castor oil or raw lined oil at the rate of one teaspoonful to one tablespoonful for very young pigs may assist in relieving scours or enteritis. In severe cases of scours or enteritis, an enteritis powder may be used very effectively. The addition to the feed twice daily of three ounces of a mixture containing four parts sodium bicarbonate, two parts of bismuth subnitrate and one part of salol for each 1,000 pounds of live weight is effective. In very young pigs a teaspoonful of this powder is used twice daily.

Round Worms. Round worms and other intestinal worms are common causes of diseases of hogs. Round worms are easily detected in the intestines of affected pigs as round yellowish-white worms, eight to twelve inches long. These worms are present on practically every farm where hogs have been grown for a number of years. This worm (*Ascaris suis*) lays as many as 100,000 eggs annually that pass out with the feces into the hog lot. These eggs embryonate or become infectious after one or two weeks under favorable conditions. When taken into the digestive tract of young pigs with feed or water, they hatch out and the young worms burrow through the intestine into the blood stream and pass to the lungs




Fig. 23.—Where Round worms are found other internal parasites and filth-borne diseases are also present. Sanitary rearing is the only permanent solution of these parasite problems. *Courtesy Bureau of Animal Industry, U. S. A.*

and other organs. They stay there for about a week or ten days when they migrate up or are coughed up the windpipe (*trachea*) into the mouth. They are then swallowed by the pig and develop into mature worms in the small intestines. About six or eight weeks is required for this process. (Fig. 23). A few round worms are not necessarily the sole cause of a pig becoming a runt, but the fact that a pig has round worms indicates that it has been grown under conditions which favor the development of other disease con-

ditions such as lung worms, coccidiosis, and other filth-borne diseases, all of which are detrimental and which when combined are devastating to the health of the pig.

Several treatments for round worms have been developed. They are fairly effective in removing round worms in four to five months old pigs if they are administered carefully and to each individual separately (*Fig. 24*). These treatments are not effective against lung worms, coccidiosis, or intestinal infection commonly associated with round worms, nor do they repair the damage already done to the intestinal tract of the pigs.

The most satisfactory treatment for the round worm is as follows:

1. Fast animals for 18 hours.
2. Treat with individual doses using a dose syringe.
3. Mix one ounce oil of chenopodium with one pound of castor oil or one-half pound oil of chenopodium with one gallon of castor oil.
4. Give one ounce (two tablespoonfuls) of the mixture to 50-pound pigs and two ounces for 100-pound pigs. For larger animals use more castor oil up to three or four ounces, but do not increase the oil of chenopodium.




Fig. 24.—Where pills or capsules are to be given hogs, the mouth speculum and balling gun are great time savers.

The use of tetra-chlorethylene in the treatment of hogs for worms is somewhat less effective than the above, but it is used quite generally. This drug is sold under trade names in two and five cubic centimeter capsules with directions. It should be given after 18 hours fasting and should be followed in four hours by a laxative.

The feeding of slaked coal, copperas, lye, blue vitrol or sulphur is not recommended as a means of controlling or preventing round worms in pigs. None of these substances has proved of value.

The complete prevention of worm infestation by the use of the hog lot sanitation already suggested is the only safe and sane way to combat worms.

Coccidiosis. Coccidiosis of pigs has recently been demonstrated to be a very common disease of young and partially matured pigs. The organism responsible for this condition is not visible to the naked eye. It is found in the intestinal tract and produces irritation and even bleeding by burrowing into and rupturing the cells lining the intestine.

Coccidiosis of pigs is associated with worms and intestinal infections causing scours and dysentery of young pigs. Affected pigs have poor appetites, are small and do not grow. Many young pigs die with an acute diarrhea that may become bloody. Less severely affected pigs simply become runts or may die later from infections that gain a foothold during the attack of coccidiosis.

Coccidiosis may be diagnosed best by a microscopic examination of scrapings from the intestines of an affected pig, or less satisfactorily by examination of the feces. The practical procedure for a grower to follow in controlling this disease is to clean out the bowels with a laxative and place the affected pigs in a dry place where all fecal matter can be removed at weekly intervals. They may be shifted to a clean place on a dry pasture at weekly intervals instead, if this seems more practical.

An important point to remember in controlling these intestinal parasites and filth-borne diseases is that the organisms or eggs responsible for them are all discharged with the feces and contaminate anything with which the feces come in contact. Fecal contaminated material or environment is the bane of hog production.

Hog Lice. Hog lice occur most often back of the ears or in localities that are protected and where the skin is thin and tender. Both hog lice and hog mange can be controlled by dipping the affected animals in crude oil. Farmers' Bulletin No. 1085 of the U. S. Department of Agriculture, "Hog Lice and Hog Mange, Methods of Control and Eradication", gives methods of dipping, hand treatment and other methods of application of crude oil, old crank case oil or the use of coal-tar-cresote dips in the control of lice and mange of hogs. Dipping with crude oil is the most satisfactory. Where a dipping vat is not available, hogs may be treated with a spray pump oiler. The treated hogs should not be confined too closely indoors during warm weather.

Rickets. Rickets in pigs is caused by a deficiency of calcium or other bone forming materials in the body. Pigs that are confined indoors on a ration low in calcium suffer most from this condition. Young pigs with rickets have sore feet, crooked legs and poor appetites. Pigs that are being grown on exclusive grain rations many times go down in the rear quarters and are "creepy". This is a symptom of rickets or weakness of the bones. Supplying pigs with bone meal, tankage, limestone or other calcium-carrying minerals together with plenty of sunshine or vitamin D supplements such as alfalfa leaves or cod liver oil, will prevent this condition. Pigs fed a ration containing liberal amounts of alfalfa or pigs on pasture do not ordinarily suffer from rickets. Rickets is common in Idaho

where grains are fed exclusively. It is not common where alfalfa is produced abundantly and fed extensively to the hogs (*See Minerals*).

Goitre. Goitre (hairlessness) in pigs is caused by a deficiency of iodine in the ration of the sow during the gestation period. The deficiency of iodine which causes enlargement of the thyroid gland is evidenced by large necks on the young pigs and by the partial or practically complete absence of hair over the entire body. Such pigs, if not still-born, are weak and die soon after birth. This condition may be prevented by adding one ounce of potassium iodide to each 100 pounds of the salt mixture used in the ration of the sow (*See Hairless Pigs*).

Anemia. Anemia is most often found in early-spring pigs that are farrowed in a central hog house and that do not have access to feed other than the sow's milk for the first two weeks of life. Sow's milk is deficient in iron. Pigs consequently suffer from a lack of red blood cell forming material and many die of anemia before they are two weeks of age. Affected pigs are weak and extremely short of breath. They may appear in excellent flesh—even over fat. The slightest exertion causes them to thump or breathe rapidly. Less severely affected pigs become runts and many die later in life from other causes.

Anemia can be prevented by dosing the pigs with iron sulphate and copper sulphate. A mixture containing seventeen parts of iron sulphate and three parts of copper sulphate used at the rate of four ounces to a quart of water may be applied to the udder of the sow twice weekly. This application will normally supply enough iron for the suckling pigs to prevent anemia. It has recently been proved that the liberal feeding of clean soil or sod to young pigs confined indoors aids materially in preventing the development of anemia.

Thumps. Thumps is a symptom of some disturbance of respiration. Pigs showing this symptom breathe with difficulty, often rapidly and spasmodically. The disease most often responsible for the development of thumps is anemia (*See Anemia*). Pigs suffering from pneumonia often show the same symptoms. The common round worm is often responsible for pneumonia since it passes through the lungs in its development (*See Round Worms*). Lung worms may also be responsible for this disease.

Piles in Pigs. Piles in pigs is somewhat similar to hemeroids or piles in humans. It is a protrusion of the wall of the rectum past the natural body opening. It is brought on by any feed or condition that tends to develop constipation. It is best relieved or prevented by adding bran, alfalfa meal, or other laxative material to the ration. If the protruding bowel remains out for several days, it may be safely removed by cutting it off close to the skin with a sharp knife.

Hernia or Rupture. Umbilical and scrotal hernias are most common. The passage of the intestines through openings of the abdominal wall into sacks under the skin of the belly or into the scrotum produces a hernia. Pigs with such defects should not be used for breeding stock as these conditions are heritable. Umbilical hernias ordinarily do not need to be reduced. Scrotal hernias do need

to be reduced at the time of castration to prevent the bowels from escaping when the sack is opened. This may be done by cutting through the scrotum down to the sack or tunic covering the testicle, twisting it up over the testicle until the hernia material is forced back into the abdomen. The sack is then tied off as high up as possible and the sack and testicles are removed below the ligature with a knife.

Abscesses. Abscesses may develop at any point on the body. They usually occur just under the skin and are most often seen around the jaw or along the throat. When they have headed up, that is, become soft at one point, they should be lanced and irrigated with a mild antiseptic solution.

Precautions for Use of Drugs in Feed. Where drugs are used in the feed, care should be taken to segregate the animals into comparable groups for size, age and degree of sickness, using an amount of feed that the respective groups will consume. The more vigorous and healthy hogs would otherwise obtain more feed and consequently more of the medicine, thus limiting the amount animals more seriously affected and most needing treatment would get.

Restraint of Hogs. Pass a slip loop on the end of a rope over the upper jaw back of the tusks, tighten it up and snub to a convenient post or fence. Any hog, regardless of its size, so confined will simply pull back and can be drenched or otherwise treated or handled conveniently.

Summary

Purposely, very little has been said in this Bulletin on methods of treatment of swine diseases. Sanitation has been emphasized because most of the hog disease problems in this state are due to the lack of sanitation. Disease conditions developing from lax sanitation are, furthermore, not amenable to treatment. Prevention is, therefore, the only method that will satisfactorily solve the swine disease problem.

Swine sanitation includes placing a clean sow in a clean pen to farrow, moving her and the pigs to clean pasture and maintaining cleanliness in feeding and management methods while on pasture. The pigs must be prevented from coming in contact with old hog lots, and runs or lanes that are contaminated with hog manure. Such a procedure will make it possible to raise more pigs per litter; will reduce the number of runts; increase the rate of gain; and have the pigs ready for market earlier and at less expense.